



ELSEVIER

Contents lists available at SciVerse ScienceDirect

Journal of Combinatorial Theory, Series A

www.elsevier.com/locate/jcta


A spectral excess theorem for nonregular graphs

Guang-Siang Lee¹, Chih-wen Weng¹

Department of Applied Mathematics, National Chiao Tung University, 1001 Ta Hsueh Road, Hsinchu, Taiwan, ROC

ARTICLE INFO

Article history:

Received 27 September 2011

Available online xxxx

Keywords:

Distance-regular graphs

Eigenvalues

Spectral excess theorem

ABSTRACT

The spectral excess theorem asserts that the average excess is, at most, the spectral excess in a regular graph, and equality holds if and only if the graph is distance-regular. An example demonstrates that this theorem cannot directly apply to nonregular graphs. This paper defines average weighted excess and generalized spectral excess as generalizations of average excess and spectral excess, respectively, in nonregular graphs, and proves that for any graph the average weighted excess is at most the generalized spectral excess. Aside from distance-regular graphs, additional graphs obtain the new equality. We show that a graph is distance-regular if and only if the new equality holds and the diameter D equals the spectral diameter d . For application, we demonstrate that a graph with odd-girth $2d + 1$ must be distance-regular, generalizing a recent result of van Dam and Haemers.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Throughout this paper, let $G = (VG, EG)$ be a connected graph on n vertices, with diameter D , adjacency matrix A , and distance function ∂ . Assume that A has $d + 1$ distinct eigenvalues $\lambda_0 > \lambda_1 > \dots > \lambda_d$ with corresponding multiplicities $m_0 = 1, m_1, \dots, m_d$. From the spectrum of G we then define an inner product $\langle \cdot, \cdot \rangle_\Delta$ on the vector space $\mathbb{R}_d[x]$ of real polynomials of degree at most d . It is well known that $\mathbb{R}_d[x]$ has a unique orthogonal basis $p_0(x), p_1(x), \dots, p_d(x)$, satisfying $\deg p_i(x) = i$ and $\langle p_i(x), p_i(x) \rangle_\Delta = p_i(\lambda_0)$ for $0 \leq i \leq d$. The number $\bar{k}_d := |\{(u, v) \mid u, v \in VG, \partial(u, v) = d\}|/n$ is called the *average excess* of G , and the number $p_d(\lambda_0)$ is called the *spectral excess* of G . The spectral excess theorem, proposed by Fiol and Garriga [9], states that

$$\bar{k}_d \leq p_d(\lambda_0) \quad (1)$$

E-mail addresses: tremolo.am96g@nctu.edu.tw (G.-S. Lee), weng@math.nctu.edu.tw (C.-w. Weng).

¹ Fax: +886 3 5724679.

if G is regular; and equality holds if and only if G is distance-regular. For short proofs, see [5,11]. Furthermore, see [4,3] for some generalizations.

Example 2.2 indicates that (1) cannot directly apply to nonregular graphs. In addition, (1) is trivial if $D < d$, because $\bar{k}_d = 0$. Consequently, we provide a generalization of the spectral excess theorem to make it applicable to nonregular graphs using a global approach. Fiol, Garriga, and Yebra [13] also considered nonregular graphs, using a local approach, however. In Section 3 we define the *average weighted excess* δ_D and the *generalized spectral excess* $p_{\geq D}(\lambda_0) := p_D(\lambda_0) + p_{D+1}(\lambda_0) + \dots + p_d(\lambda_0)$, as generalizations of average excess and spectral excess, respectively, in nonregular graphs, and prove that

$$\delta_D \leq p_{\geq D}(\lambda_0). \tag{2}$$

As the spectral excess theorem, distance-regular graphs also attain equality in (2). Moreover, additional graphs obtain equality (see Remark 3.3 for more details). Furthermore, under the assumption $D = d$, we show in Theorem 3.5 that G is distance-regular if and only if equality in (2) holds.

The *odd-girth* of a graph is the length of its shortest odd cycle. For application, in Section 4, we demonstrate that a graph with odd-girth $2d + 1$ must be distance-regular. The same consequence was previously proved by van Dam and Haemers [6] under the assumption that G is regular. Moreover, our result is the solution of a problem raised in the same paper (though these authors also proved this result for the case $d + 1 = 3$ and claimed to have proofs for the cases $d + 1 \in \{4, 5\}$). Because the odd-girth is determined according to the spectrum, this result is also a generalization of the spectral characterization of the generalized odd graphs [15,16].

2. Preliminaries

In this section, we review the concept of orthogonal polynomials related to G . The basic idea is to generalize the study of distance-regular graphs (see [1,2,18]). The *spectrum* of G is denoted by the multiset $\text{sp } G = \{\lambda_0^{m_0}, \lambda_1^{m_1}, \dots, \lambda_d^{m_d}\}$, and the parameter d is called the *spectral diameter* of G . It is well known that $D \leq d$ and $Z(x) := \prod_{i=0}^d (x - \lambda_i)$ is the *minimal polynomial* of G [1, Chapter 2]. From the spectrum $\text{sp } G = \{\lambda_0^{m_0}, \lambda_1^{m_1}, \dots, \lambda_d^{m_d}\}$ we consider the $(d + 1)$ -dimensional vector space $\mathbb{R}_d[x] \cong \mathbb{R}[x]/(Z(x))$ of real polynomials of degree at most d with inner product defined by

$$\langle p(x), q(x) \rangle_{\Delta} := \sum_{i=0}^d \frac{m_i}{n} p(\lambda_i)q(\lambda_i) = \text{tr}(p(A)q(A))/n.$$

Using this inner product, there is a unique system of orthogonal polynomials $p_i(x)$, $i = 0, 1, \dots, d$, where $\deg p_i(x) = i$ and $\langle p_i(x), p_j(x) \rangle_{\Delta} = p_i(\lambda_0)$ [17]. These polynomials $p_0(x), p_1(x), \dots, p_d(x)$ are referred to as the *predistance polynomials* of G . Moreover, the sum of all predistance polynomials gives the *Hoffman polynomial* $H(x)$ [14]:

$$H(x) := n \prod_{i=1}^d \frac{x - \lambda_i}{\lambda_0 - \lambda_i} = p_0(x) + p_1(x) + \dots + p_d(x), \tag{3}$$

no matter whether the graph G is regular or not. For a proof, see for instance [5]. Let α be the eigenvector of A corresponding to λ_0 such that $\alpha^t \alpha = n$ and all entries of α are positive. Notice that $\alpha = (1, 1, \dots, 1)^t$ if and only if G is regular. The following lemma generalizes [14] to nonregular graphs, which was considered in [8, p. 117].

Lemma 2.1. *Let G be a graph with adjacency matrix A . Then, $H(A) = \alpha \alpha^t$. Moreover, G is regular if and only if $H(A) = J$, the all 1's matrix. \square*

Recall that the spectral excess theorem states that if G is regular then (1) holds. The following example demonstrates that the regularity assumption of G is necessary.

Example 2.2. Let P_3 be a path of three vertices, with spectrum $\text{sp } P_3 = \{\sqrt{2}, 0, -\sqrt{2}\}$. One can check that $p_0(x) = 1$, $p_1(x) = 3\sqrt{2}x/4$, and $p_2(x) = 3(x^2 - 4/3)/4$. Notice that $k_2 = 2/3$ and $p_2(\lambda_0) = 1/2$. This shows that (1) does not hold.

3. A weighted spectral excess theorem

Let A_i be the i -th distance matrix of G , i.e., the $n \times n$ matrix with rows and columns indexed by the vertex set VG such that

$$(A_i)_{uv} = \begin{cases} 1 & \text{if } \partial(u, v) = i, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, $A_0 = I$ is the identity matrix and $A_1 = A$ is the adjacency matrix. Define $\tilde{A}_i := A_i \circ H(A)$, where “ \circ ” is the entrywise product of matrices. Notice that \tilde{A}_i can be regarded as a “weighted” version of A_i since

$$(\tilde{A}_i)_{uv} = \begin{cases} \alpha_u \alpha_v & \text{if } \partial(u, v) = i, \\ 0 & \text{otherwise} \end{cases}$$

by Lemma 2.1. This approach of giving weights, the entries of the positive eigenvector, to the vertices of a nonregular graph has been recently used many times in the literature (see, for instance, [12,9,10, 7]). For instance, this idea was crucial to introduce the concept of pseudo-distance-regularity and the corresponding (u -local) i -distance matrices in [12]. The reason that \tilde{A}_i is weighted in the above way is to have the property that

$$\tilde{A}_0 + \tilde{A}_1 + \dots + \tilde{A}_D = H(A) = p_0(A) + p_1(A) + \dots + p_d(A)$$

by Lemma 2.1 and (3). Define $\delta_i := \langle \tilde{A}_i, \tilde{A}_i \rangle$ and $p_{\geq D}(x) := p_D(x) + p_{D+1}(x) + \dots + p_d(x)$. The number δ_D is referred to as the *average weighted excess* and $p_{\geq D}(\lambda_0)$ as the *generalized spectral excess* of G . Notice that if $D = d$ then $p_{\geq D}(x) = p_d(x)$. For any two $n \times n$ real symmetric matrices M and N , define the inner product

$$\langle M, N \rangle := \frac{1}{n} \text{tr}(MN) = \frac{1}{n} \sum_{i,j} (M \circ N)_{ij}.$$

Moreover, let

$$\text{Proj}_N(M) := \frac{\langle N, M \rangle}{\langle N, N \rangle} N$$

denote the projection of M onto $\text{Span}\{N\}$. The proof of the following lemma is basically the same as in [11, Lemma 1] (the only difference is the use of “weights”, the entries of α , on the vertices of G).

Lemma 3.1. $\text{Proj}_{\tilde{A}_D}(p_{\geq D}(A)) = \tilde{A}_D$. \square

From Lemma 3.1, we immediately have the following theorem.

Theorem 3.2. *Let G be a connected graph with diameter D . Then $\delta_D \leq p_{\geq D}(\lambda_0)$ with equality if and only if $\tilde{A}_D = p_{\geq D}(A)$.* \square

Revisiting the case when G is the path on three vertices P_3 described in Example 2.2. Notice that $D = d = 2$, $\alpha = (\sqrt{3}/2, \sqrt{6}/2, \sqrt{3}/2)^t$ and

$$\tilde{A}_D = \begin{pmatrix} 0 & 0 & 3/4 \\ 0 & 0 & 0 \\ 3/4 & 0 & 0 \end{pmatrix}.$$

Then $\delta_D = 3/8 \leq 1/2 = p_{\geq D}(\lambda_0)$ satisfies inequality in Theorem 3.2.

Remark 3.3. If G is regular with diameter $D = 2$, then equality in Theorem 3.2 holds. Indeed, $\tilde{A}_2 = A_2 = J - I - A = H(A) - I - A = p_{\geq 2}(A)$.

The graph described in Remark 3.3 is a special case of *distance-polynomial graphs* [19]. It would be interesting to characterize the graphs which satisfy equality in Theorem 3.2. We complete this characterization under the assumption $D = d$ in Theorem 3.5. Since $p_0(A) = I$, the following result is simple, but plays a crucial role in proving the regularity of a graph.

Lemma 3.4. $\tilde{A}_0 = p_0(A)$ if and only if G is regular. \square

The following theorem is a “weighted” version of [11, Proposition 2]. The proof is essentially the same except that here the use of weights on vertices is taken into account.

Theorem 3.5. Let G be a connected graph of diameter D which is equal to the spectral diameter d . Then $\tilde{A}_D = p_D(A)$ if and only if $\tilde{A}_i = p_i(A)$ for $0 \leq i \leq D - 1$. Moreover, if $\tilde{A}_D = p_D(A)$ then G is distance-regular.

Proof. The first part of the proof is essentially the same as in [11, Proposition 2]. Suppose $\tilde{A}_D = p_D(A)$. By the first part we conclude that $\tilde{A}_0 = p_0(A)$, and thus G is regular by Lemma 3.4. Hence G is distance-regular by [11, Proposition 2]. \square

4. Graphs with odd-girth $2d + 1$

In this section, assume that G has odd-girth $2d + 1$. For application of Theorem 3.5, we demonstrate that G has diameter $D = d$ and G must be distance-regular. The proof is basically identical to [6, pp. 487–488], except for the weights on the vertices. Let $c = n / \prod_{i=1}^d (\lambda_0 - \lambda_i)$, which is the leading coefficient of the Hoffman polynomial $H(x)$. Notice that the trace $\text{tr}(A^{2d+1})$ of A^{2d+1} is nonzero since G has odd-girth $2d + 1$. The following lemma determines the average weighted excess δ_D and the generalized spectral excess $p_{\geq D}(\lambda_0)$.

Lemma 4.1. (See [6].) $\delta_D = c^2 \text{tr}(A^{2d+1}) / (n \sum_{i=0}^d \lambda_i) = p_d(\lambda_0)$. In particular, $D = d$. \square

From Lemma 4.1 and Theorem 3.5, we immediately have the following theorem.

Theorem 4.2. A connected graph with $d + 1$ distinct eigenvalues and odd-girth $2d + 1$ is distance-regular. \square

Acknowledgments

The authors are grateful to the anonymous referees for careful reading and valuable suggestions that improved the presentation. This research is supported by the National Science Council of Taiwan, ROC under the project NSC 99-2115-M-009-005-MY3.

References

- [1] N. Biggs, Algebraic Graph Theory, Cambridge University Press, Cambridge, 1993.
- [2] A.E. Brouwer, A.M. Cohen, A. Neumaier, Distance-Regular Graphs, Springer-Verlag, Berlin, 1989.
- [3] C. Dalfó, E.R. van Dam, M.A. Fiol, E. Garriga, Dual concepts of almost distance-regularity and the spectral excess theorem, Discrete Math. (2012), <http://dx.doi.org/10.1016/j.disc.2012.03.003>.
- [4] C. Dalfó, E.R. van Dam, M.A. Fiol, E. Garriga, B.L. Gorissen, On almost distance-regular graphs, J. Combin. Theory Ser. A 118 (2011) 1094–1113.
- [5] E.R. van Dam, The spectral excess theorem for distance-regular graphs: a global (over)view, Electron. J. Combin. 15 (1) (2008) R129.
- [6] E.R. van Dam, W.H. Haemers, An odd characterization of the generalized odd graphs, J. Combin. Theory Ser. B 101 (2011) 486–489.
- [7] M.A. Fiol, Eigenvalue interlacing and weight parameters of graphs, Linear Algebra Appl. 290 (1999) 275–301.

- [8] M.A. Fiol, Algebraic characterizations of distance-regular graphs, *Discrete Math.* 246 (2002) 111–129.
- [9] M.A. Fiol, E. Garriga, From local adjacency polynomials to local pseudo-distance-regular graphs, *J. Combin. Theory Ser. B* 71 (1997) 162–183.
- [10] M.A. Fiol, E. Garriga, On the algebraic theory of pseudo-distance-regularity around a set, *Linear Algebra Appl.* 298 (1999) 115–141.
- [11] M.A. Fiol, S. Gago, E. Garriga, A simple proof of the spectral excess theorem for distance-regular graphs, *Linear Algebra Appl.* 432 (2010) 2418–2422.
- [12] M.A. Fiol, E. Garriga, J.L.A. Yebra, On a class of polynomials and its relation with the spectra and diameters of graphs, *J. Combin. Theory Ser. B* 67 (1996) 48–61.
- [13] M.A. Fiol, E. Garriga, J.L.A. Yebra, Locally pseudo-distance-regular graphs, *J. Combin. Theory Ser. B* 68 (1996) 179–205.
- [14] A.J. Hoffman, On the polynomial of a graph, *Amer. Math. Monthly* 70 (1963) 30–36.
- [15] T. Huang, Spectral characterization of odd graphs O_k , $k \leq 6$, *Graphs Combin.* 10 (1994) 235–240.
- [16] T. Huang, C. Liu, Spectral characterization of some generalized odd graphs, *Graphs Combin.* 15 (1999) 195–209.
- [17] G. Szegő, *Orthogonal Polynomials*, fourth edition, Amer. Math. Soc. Colloq. Publ., vol. XXIII, American Mathematical Society, Providence, RI, 1975.
- [18] P. Terwilliger, The subconstituent algebra of an association scheme I, *J. Algebraic Combin.* 1 (1992) 363–388.
- [19] P.M. Weichsel, On distance-regularity in graphs, *J. Combin. Theory Ser. B* 32 (1982) 156–161.