

DETERMINING COEFFICIENTS FOR A FRACTIONAL p -LAPLACE EQUATION FROM EXTERIOR MEASUREMENTS

MANAS KAR, YI-HSUAN LIN, AND PHILIPP ZIMMERMANN

ABSTRACT. We consider an inverse problem of determining the coefficients of a fractional p -Laplace equation in the exterior domain. Assuming suitable local regularity of the coefficients in the exterior domain, we offer an explicit reconstruction formula in the region where the exterior measurements are performed. This formula is then used to establish a *global uniqueness* result for real-analytic coefficients. In addition, we also derive a stability estimate for the unique determination of the coefficients in the exterior measurement set.

Keywords. Inverse problems, exterior determination, fractional gradient, fractional divergence, fractional p -Laplacian.

Mathematics Subject Classification (2020): Primary 35R30; secondary 26A33, 42B37, 46F12

CONTENTS

1. Introduction	1
1.1. The p -Calderón problem	2
1.2. Nonlocal inverse problems	3
1.3. Main results	4
1.4. Organization of the article	4
2. Preliminaries	5
2.1. Function spaces	5
2.2. Nonlocal operators	6
Conventions	7
3. The forward problem and DN map	7
3.1. Auxiliary lemma	7
3.2. Well-posedness	7
3.3. DN maps	11
4. Exterior reconstruction	12
5. Proofs of Proposition 1.2, 1.3 and 1.4	19
Acknowledgments	19
References	19

1. INTRODUCTION

In this article, we study an inverse problem for a fractional p -Laplace equation. To formulate the problem, let us consider a partial differential equation (PDE) of the form

$$(1.1) \quad \operatorname{Div}_s (\sigma |d_s u|^{p-2} d_s u) = 0,$$

where $1 < p < \infty$, $\sigma = \sigma(x, y): \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the uniform ellipticity condition

$$(1.2) \quad \lambda \leq \sigma(x, y) \leq \lambda^{-1}, \text{ for all } x, y \in \mathbb{R}^n,$$

and for some constant $\lambda > 0$. Later we call this operator appearing in (1.1) *weighted fractional p -Laplacian*. Here, $d_s u$ denotes the *fractional s -gradient* and Div_s is its adjoint, which is the *fractional s -divergence*, with respect to the measure

$$d\mu = \frac{dxdy}{|x-y|^n} \text{ on } \mathbb{R}^n \times \mathbb{R}^n.$$

More concretely, for any function $u: \mathbb{R}^n \rightarrow \mathbb{R}$ and $s \in (0, 1)$, the fractional s -gradient and s -divergence could be defined by

$$d_s u(x, y) = \frac{u(x) - u(y)}{|x - y|^s} \quad \text{and} \quad \langle \text{Div}_s u, \varphi \rangle = \int_{\mathbb{R}^{2n}} \frac{u(x, y) d_s \varphi(x, y)}{|x - y|^n} dxdy,$$

respectively, for $x, y \in \mathbb{R}^n$ and for all $\varphi \in C_c^\infty(\mathbb{R}^n)$ (see for example [MS18] and Section 2 for detailed definitions). The weighted fractional p -Laplacian is weakly defined by

$$\begin{aligned} & \langle \text{Div}_s(\sigma |d_s u|^{p-2} d_s u), \varphi \rangle \\ & := \int_{\mathbb{R}^{2n}} \sigma(x, y) |u(x) - u(y)|^{p-2} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+sp}} dxdy, \end{aligned}$$

for any $\varphi \in C_c^\infty(\mathbb{R}^n)$.

Next, we are going to formulate an inverse problem related to (1.1). The observations of our inverse problem are encoded in the *Dirichlet-to-Neumann* (DN) map, which is formally defined by

$$\Lambda_\sigma(f) = \text{Div}_s(\sigma |d_s u|^{p-2} d_s u) \Big|_{\Omega_e}$$

where $\Omega_e := \mathbb{R}^n \setminus \bar{\Omega}$ denotes the exterior domain and u_f is the unique solution of

$$(1.3) \quad \begin{cases} \text{Div}_s(\sigma |d_s u|^{p-2} d_s u) = 0 & \text{in } \Omega, \\ u = f & \text{in } \Omega_e. \end{cases}$$

Here we have assumed the well-posedness of (1.1) at the moment (the proof will be given in Section 3) and that the weighted fractional p -Laplacian of u induces at least a distribution on Ω_e . Then we ask the following question:

Question 1. *Let $W \subset \Omega_e$ be a given nonempty open set and assume that the coefficients σ_1, σ_2 satisfy $\Lambda_{\sigma_1} f|_W = \Lambda_{\sigma_2} f|_W$, for all $f \in C_c^\infty(W)$. If $\Sigma_j: \mathbb{R}^n \rightarrow \mathbb{R}$ is given by $\Sigma_j(x) = \sigma_j(x, x)$ for $j = 1, 2$, can we conclude $\Sigma_1 = \Sigma_2$ in W ?*

In the special case of coefficients of the form $\sigma(x, y) = \gamma^{1/2}(x)\gamma^{1/2}(y)$ this generalizes recent results for the fractional conductivity equation (i.e. $p = 2$) by the last author (see [RZ22b, RZ22a, CRZ22, RZ22c, CRTZ22] for the elliptic and [LRZ22] for the parabolic case).

As $s = 1$, the related inverse problem to the equation

$$(1.4) \quad \text{div}(\gamma |\nabla u|^{p-2} \nabla u) = 0$$

is the so called (classical) p -Calderón problem, where $\gamma: \mathbb{R}^n \rightarrow \mathbb{R}$ is a positive scalar function¹, and we next discuss about it.

1.1. The p -Calderón problem. In the p -Calderón problem the DN map is strongly given by

$$f \mapsto \Lambda_\gamma^p f = \gamma |\nabla u_f|^{p-2} \partial_\nu u_f \Big|_{\partial\Omega},$$

where u_f is the unique solution to

$$(1.5) \quad \begin{cases} \text{div}(\gamma |\nabla u|^{p-2} \nabla u) = 0 & \text{in } \Omega, \\ u = f & \text{on } \partial\Omega. \end{cases}$$

¹In this paper, we use the notation $\gamma = \gamma(x): \mathbb{R}^n \rightarrow \mathbb{R}$ and $\sigma = \sigma(x, y): \mathbb{R}^{2n} \rightarrow \mathbb{R}$.

Now the inverse problem is to ask whether one can determine the coefficient γ uniquely from the knowledge of the (nonlinear) DN map Λ_γ^p ? Note that in the special case $p = 2$ this reduces to the classical Calderón problem (see [Cal06, KV84, SU87]).

Moreover, if $\gamma = 1$, the partial differential operator in (1.5) becomes the p -Laplacian $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$, which appears in the study of nonlinear dielectrics [GK03, TW94a, TW94b, LK98], plastic moulding [Aro96], nonlinear fluids [AR06, AJ92, GR03, Idi08] and others. In [SZ12] the authors proved by using p -harmonic functions (i.e. functions solving (1.4)) introduced by Wolff [Wol07] that the nonlinear DN map Λ_γ^p determines uniquely γ on the boundary. Later, Brander showed in [Bra16] that the DN map also determines the normal derivative $\partial_\nu \gamma$ on $\partial\Omega$. These results can be seen as a zeroth and first order analogue of the boundary determination result of Kohn and Vogelius [KV84] for the Calderón problem. Since not all coefficients of the Taylor series around a boundary point, as in the classical Calderón problem, are known from the DN map Λ_γ^p , it cannot be used to determine real-analytic coefficients in the interior of Ω . Meanwhile, the authors [BHKS18, GKS16, BKS15, KW21, BIK18] studied inverse problems for (weighted) p -Laplace equations by utilizing monotonicity methods. Recently, in the work [CF24], the authors recover the leading coefficient for the weighted p -Laplace equation in the plane, without assuming additional conditions for the leading coefficient, and they also showed another uniqueness result in the dimension 3 and higher with some suitable assumptions.

1.2. Nonlocal inverse problems. In recent years, many different Calderón type inverse problems for nonlocal operators have been studied. The prototypical example is the inverse problem for the fractional Schrödinger operator $(-\Delta)^s + q$ with $q \in L^\infty(\Omega)$, which was first considered in [GSU20] and initiated many of the later developments. The main ingredients in solving this Calderón problem are an *Alessandrini identity*, the *UCP* (unique continuation property) and the closely related *Runge approximation*. It is worth noticing that the UCP and the approximation are much stronger in the nonlocal case than in the local ones, which is mainly possible, because solutions to $(-\Delta)^s + q$ are much less rigid than the ones to the local Schrödinger equation $-\Delta + q$. By using a similar approach, one can solve a variety of inverse problems for nonlocal operators whose corresponding local counterpart is still open. For further details we refer to the works [BGU21, CMR21, CMRU22, GLX17, CL19, CLL19, CRZ22, CLR20, FGKU21, HL19, HL20, GRSU20, GU21, Gho22, Lin22, Lin23, LL22a, LL22b, LLR20, LLU22, KIW22, RS20, RS18, RZ22a, RZ22c, RZ22b]. We point out that most of these works consider nonlocal inverse problems in which one recovers lower order coefficients instead of principal order like in the classical Calderón problem. On the other hand, in the articles [GU21, LLU22, RZ22c, RZ22b, RZ22a] the authors study nonlocal inverse problems where one is interested in determining leading order coefficients and hence they can be seen as full nonlocal analogues of the classical Calderón problem.

Let us mention that in all the previous inverse problems the leading order operator of the underlying nonlocal PDEs is linear. A first step into the direction of considering nonlinear nonlocal leading order operators was taken in the work [KRZ22] by the first and the last author. A crucial advantage of the operators L studied in this work is that they have the UCP, that is, if $u: \mathbb{R}^n \rightarrow \mathbb{R}$ is a sufficiently regular function and $Lu = u = 0$ in an open set $V \subset \mathbb{R}^n$, then $u \equiv 0$ in \mathbb{R}^n . But in contrast the UCP for the operators in (1.4) is only known to hold in $n = 2$ dimensions but for $n \geq 3$ it is a difficult open problem. Similarly, it is not known whether the operators in (1.1) have the UCP.

1.3. Main results. The main theorem of this article is the following exterior reconstruction result on the diagonal extending [CRZ22, Proposition 1.5].

Theorem 1.1 (Exterior reconstruction on the diagonal). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $W \subset \Omega_e$ a nonempty open set, $0 < s < 1$, $1 < p < \infty$ and $x_0 \in W \Subset \Omega_e$. Then there exists a sequence $(\Phi_N)_{N \in \mathbb{N}} \subset C_c^\infty(W)$ such that*

- (i) *for all $N \in \mathbb{N}$ it holds $[\Phi_N]_{W^{s,p}(\mathbb{R}^n)} = 1$,*
- (ii) *for all $0 \leq t < s$ there holds $\|\Phi_N\|_{W^{t,p}(\mathbb{R}^n)} \rightarrow 0$ as $N \rightarrow \infty$*
- (iii) *and $\text{supp}(\Phi_N) \rightarrow \{x_0\}$ as $N \rightarrow \infty$.*

Moreover, if $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies uniformly elliptic condition (1.2) such that $(\sigma(x, \cdot): W \rightarrow \mathbb{R})_{x \in \mathbb{R}^n}$ is equicontinuous at x_0 and $\sigma(\cdot, x_0) \in C(W)$, then there holds

$$(1.6) \quad \sigma(x_0, x_0) = \lim_{N \rightarrow \infty} \langle \Lambda_\sigma \Phi_N, \Phi_N \rangle.$$

As an immediate consequence of the formula (1.6), we obtain the following results on exterior determination, exterior stability and global uniqueness for real-analytic coefficients.

Proposition 1.2 (Exterior determination on the diagonal). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $W \subset \Omega_e$ be a nonempty open set, $0 < s < 1$ and $1 < p < \infty$. Assume that $\sigma_j: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the conditions of Theorem 1.1 for $j = 1, 2$ and set $\Sigma_j(x) := \sigma_j(x, x)$ for $x \in \mathbb{R}^n$, $j = 1, 2$. Suppose $\Lambda_{\sigma_1} f|_W = \Lambda_{\sigma_2} f|_W$, for all $f \in C_c^\infty(W)$, then there holds $\Sigma_1(x) = \Sigma_2(x)$ for all $x \in W$.*

Proposition 1.3 (Exterior stability on the diagonal). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $W \subset \Omega_e$ a nonempty open set, $0 < s < 1$ and $1 < p < \infty$. Assume that $\sigma_j: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the conditions of Theorem 1.1 for $j = 1, 2$ and set $\Sigma_j(x) = \sigma_j(x, x)$ for $x \in \mathbb{R}^n$, $j = 1, 2$. Then we have*

$$\|\Sigma_1 - \Sigma_2\|_{L^\infty(W)} \leq \|\Lambda_{\sigma_1} - \Lambda_{\sigma_2}\|_{\widetilde{W}^{s,p}(W) \rightarrow (\widetilde{W}^{s,p}(W))^*}.$$

Proposition 1.4 (Global uniqueness on the diagonal). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $W \subset \Omega_e$ a nonempty open set, $0 < s < 1$ and $1 < p < \infty$. Assume that $\sigma_j: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the conditions of Theorem 1.1 for $j = 1, 2$ and set $\Sigma_j(x) = \sigma_j(x, x)$ for $x \in \mathbb{R}^n$, $j = 1, 2$. If $\Lambda_{\sigma_1} f|_W = \Lambda_{\sigma_2} f|_W$ for all $f \in C_c^\infty(W)$, and Σ_j are real-analytic for $j = 1, 2$ then $\Sigma_1 = \Sigma_2$ in \mathbb{R}^n .*

Observe that Proposition 1.4 implies several global uniqueness results when one assumes that the coefficients have a product structure. For example, if for $j = 1, 2$ the coefficients $\sigma_j(x, y)$ can be written as $\sigma_j(x, y) = F(\gamma_j(x))F(\gamma_j(y))$ for some real analytic functions $\gamma_j: \mathbb{R}^n \rightarrow \mathbb{R}$, $F: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying

- (i) γ_j is uniformly elliptic for $j = 1, 2$,
- (ii) F is injective
- (iii) and for any compact interval $[a, b] \subset \mathbb{R}_+$ there exists $c > 0$ such that $F(\xi) \geq c$ for all $\xi \in [a, b]$.

Then $\Lambda_{\sigma_1} f|_W = \Lambda_{\sigma_2} f|_W$ for all $f \in C_c^\infty(W)$ implies $\gamma_1 = \gamma_2$ in \mathbb{R}^n . As a special case one could take $F(t) = \sqrt{t}$ and recovers the global uniqueness result in [CRZ22, Theorem 1.3] for real-analytic conductivities.

1.4. Organization of the article. We first recall preliminaries related to the function spaces and nonlocal operators used throughout this work in Section 2. Afterwards in Section 3 we establish well-posedness of the exterior value problem for (1.1) and introduce the related DN map. The proof of the main result, Theorem 1.1, are divided into several steps for better readability and given in Section 4. The proofs of Proposition 1.2, 1.3 and 1.4 are given in Section 5.

2. PRELIMINARIES

Throughout this article $\Omega \subset \mathbb{R}^n$ is always a bounded open set, where $n \geq 1$ is a fixed positive integer, and $0 < s < 1$. In this section, we recall the fundamental properties of the classical fractional Sobolev spaces $W^{s,p}(\mathbb{R}^n)$ and their local analogues as well as introduce the nonlocal operators which will be used later on.

2.1. Function spaces. By $L^0(\Omega)$ we label the space of (Lebesgue) measurable functions on Ω . The classical Sobolev spaces of order $k \in \mathbb{N}$ and integrability exponent $1 \leq p \leq \infty$ are denoted by $W^{k,p}(\Omega)$ and for $k = 0$ we use the convention $W^{0,p}(\Omega) = L^p(\Omega)$. Moreover, we let $W^{s,p}(\Omega)$ stand for the fractional Sobolev spaces, when $s \in (0, 1)$ and $1 \leq p < \infty$. These spaces are also called Slobodeckij spaces or Gagliardo spaces. If $1 \leq p < \infty$ and $s \in (0, 1)$, then they are defined by

$$W^{s,p}(\Omega) := \left\{ u \in L^p(\Omega); [u]_{W^{s,p}(\Omega)} < \infty \right\},$$

where

$$[u]_{W^{s,p}(\Omega)} := \left(\int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} dx dy \right)^{1/p}$$

is the so-called Gagliardo seminorm. The fractional Sobolev spaces are naturally endowed with the norm

$$\|u\|_{W^{s,p}(\Omega)} := \left(\|u\|_{L^p(\Omega)}^p + [u]_{W^{s,p}(\Omega)}^p \right)^{1/p}.$$

The space of test functions we are going to use later in the definition of weak solutions to our PDEs is:

$$\widetilde{W}^{s,p}(\Omega) := \text{closure of } C_c^\infty(\Omega) \text{ with respect to } \|\cdot\|_{W^{s,p}(\mathbb{R}^n)}.$$

Similarly, as the classical Sobolev spaces, the spaces $W^{s,p}(\mathbb{R}^n)$ are separable for $1 \leq p < \infty$ and reflexive for $1 < p < \infty$ (see [BH22, Section 7]). Since $\widetilde{W}^{s,p}(\Omega)$ is a closed subspace of $W^{s,p}(\mathbb{R}^n)$ it has the same properties. We remark that it is known that $\widetilde{W}^{s,p}(\Omega)$ coincides with the set of all functions $u \in W^{s,p}(\mathbb{R}^n)$ such that $u = 0$ almost everywhere (a.e.) in Ω^c , when $\partial\Omega \in C^0$, and with

$$W_0^{s,p}(\Omega) := \text{closure of } C_c^\infty(\Omega) \text{ with respect to } \|\cdot\|_{W^{s,p}(\Omega)},$$

whenever $\Omega \Subset \mathbb{R}^n$ has a Lipschitz boundary (see [KLL22, Section 2]).

On these spaces $\widetilde{W}^{s,p}(\Omega)$, the following Poincaré inequality holds:

Proposition 2.1 (Poincaré inequality, [KLL22, Theorem 2.8]). *Let $\Omega \Subset \mathbb{R}^n$, $0 < s < 1$ and $1 < p < \infty$, then there exists a constant $C = C(n, s, p, \text{diam}(\Omega)) > 0$ such that*

$$(2.1) \quad \|u\|_{L^p(\mathbb{R}^n)}^p \leq C [u]_{W^{s,p}(\mathbb{R}^n)}^p$$

for all $u \in \widetilde{W}^{s,p}(\Omega)$.

Remark 2.2. *In the above theorem and from now on, we write $V \Subset W$ for two open subsets $V, W \subset \mathbb{R}^n$, if V is compactly contained in W . By the proof of [KLL22, Theorem 2.8] it follows that the optimal constant $C_* > 0$ in (2.1) satisfies $C_* \leq C_1(\text{diam}(\Omega))^{sp}$ for some $C_1 = C_1(n, s, p) > 0$. Moreover, we used here that by [KLL22, Theorem 2.8] the estimate (2.1) holds for all functions $u \in C_c^\infty(\Omega)$, but then the definition of the spaces $\widetilde{W}^{s,p}(\Omega)$ implies that by approximation it holds for all functions in this space.*

2.2. Nonlocal operators. Next we introduce the fractional s -gradient d_s , the fractional s -divergence Div_s , the fractional p -Laplacian $(-\Delta)_p^s$ and the weighted fractional p -Laplacians, which are the main object of study in this article.

For this purpose, let us denote by $L^0(\bigwedge_{od}^1 \mathbb{R}^n)$ the space of measurable off diagonal vector fields, that is, the set of all functions $F: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ which are measurable with respect to the measure $d\mu := \frac{dx dy}{|x-y|^n}$ on $\mathbb{R}^n \times \mathbb{R}^n$. We next give rigorous definitions of s -gradient and s -divergence.

Definition 2.3 (s -gradient). *For any $0 < s < 1$, the s -gradient d_s is defined by $d_s: L^0(\mathbb{R}^n) \rightarrow L^0(\bigwedge_{od}^1 \mathbb{R}^n)$ such that*

$$d_s u(x, y) := \frac{u(x) - u(y)}{|x - y|^s}.$$

Moreover, one can immediately observe that it satisfies the product rule

$$(2.2) \quad d_s(\varphi\psi)(x, y) = \varphi(x)d_s\psi(x, y) + \psi(y)d_s\varphi(x, y),$$

for a.e. $x, y \in \mathbb{R}^n$ and $\varphi, \psi: \mathbb{R}^n \rightarrow \mathbb{R}$. Moreover, we call the dual operation to the s -gradient d_s the s -divergence Div_s .

Definition 2.4 (s -divergence). *For any $0 < s < 1$, the s -divergence is the unbounded operator $\text{Div}_s: L^0(\bigwedge_{od}^1 \mathbb{R}^n) \rightarrow L^0(\mathbb{R}^n)$ given by*

$$\langle \text{Div}_s F, \varphi \rangle = \int_{\mathbb{R}^{2n}} \frac{F(x, y)d_s\varphi(x, y)}{|x - y|^n} dx dy, \text{ for all } \varphi \in C_c^\infty(\mathbb{R}^n).$$

With these definitions at our disposal, we have a canonical relation to the fractional Laplacian. In fact, there holds $\text{Div}_s \circ d_s = (-\Delta)^s$ in the sense that

$$\int_{\mathbb{R}^{2n}} \frac{d_s\varphi(x, y)d_s\psi(x, y)}{|x - y|^n} dx dy = \int_{\mathbb{R}^n} (-\Delta)^s \varphi(x) \psi(x) dx,$$

for all sufficiently regular functions $\varphi, \psi: \mathbb{R}^n \rightarrow \mathbb{R}$, where $(-\Delta)^s$ denotes the fractional Laplacian of order $s \in (0, 1)$ (up to a normalization constant). On the other hand, by the above definitions the operator $\text{Div}_s(|d_s u|^{p-2} d_s u)$ is weakly given by

$$\begin{aligned} & \langle \text{Div}_s(|d_s u|^{p-2} d_s u), \varphi \rangle \\ & := \int_{\mathbb{R}^{2n}} |u(x) - u(y)|^{p-2} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+sp}} dx dy \end{aligned}$$

for all $\varphi \in C_c^\infty(\mathbb{R}^n)$.

Hence, up to normalization, this is precisely the weak formulation of the fractional p -Laplacian. Furthermore, if the function u is sufficiently smooth then the fractional p -Laplacian can be calculated in a pointwise sense by

$$(-\Delta)_p^s u(x) = C \text{P.V.} \int_{\mathbb{R}^n} |u(x) - u(y)|^{p-2} \frac{u(x) - u(y)}{|x - y|^{n+sp}} dy$$

for some constant $C = C(n, s, p) > 0$ (see [dTGCV21]), where P.V. denotes the Cauchy principal value. For example one can take $u \in C_c^\infty(\mathbb{R}^n)$ with the additional condition $\nabla u \neq 0$ when $p \in (1, \frac{2}{2-s})$. The choice of the constant only becomes important when one wants to prove the following limiting behavior

$$\begin{cases} (-\Delta)_p^s u \rightarrow (-\Delta)^s u & \text{in } \mathbb{R}^n \text{ as } p \downarrow 2, \\ (-\Delta)_p^s u \rightarrow (-\Delta)_p u & \text{in } \mathbb{R}^n \text{ as } s \uparrow 1 \end{cases}$$

(see [dTGCV21, Section 5]). Additionally, in the recent article [BPS16] (see also [DNPV12] or [BBM01]) the authors showed that if $\Omega \Subset \mathbb{R}^n$ is a bounded Lipschitz domain then there holds

$$\lim_{s \uparrow 1} (1-s)[u]_{W^{s,p}(\mathbb{R}^n)}^p = C(n, p) \|\nabla u\|_{L^p(\Omega)}^p,$$

for some positive constant $C(n, p)$ and all $u \in W_0^{1,p}(\Omega)$. This observation is then used to show that for fixed $m \in \mathbb{N}$ and $1 < p < \infty$ the Dirichlet eigenvalues $(\lambda_{m,p}^s(\Omega))_{s \in (0,1)}$ of the fractional p -Laplacian (after a suitable renormalization) converge to a multiple of the Dirichlet eigenvalue $\lambda_{m,p}^1(\Omega)$ of the p -Laplacian as $s \uparrow 1$ and any sequence of normalized eigenfunctions $(u_{m,p}^s)_{s \in (0,1)}$ (up to a subsequence) converge to a normalized eigenfunction $u_{m,p}^1$ of the p -Laplacian as $s \uparrow 1$.

In this work, we do not pursue the limit behaviour for either $s \uparrow 1$ or $p \rightarrow 2$, but we will focus on the exterior determination results for the fractional p -Laplace equation (1.3).

Furthermore, let us point out that there is also a Caffarelli–Silvestre extension type result for the fractional p -Laplacian, when one replaces the weight y^{1-2s} by y^{1-sp} (see [dTGCV21, Section 3]). Since in this result the function u is required to be C^2 regular, the authors do not see an immediate way to generalize to proof of the UCP for the fractional Laplacian in [GSU20] (see also [KRZ22]) to the fractional p -Laplacian. Finally, note that if the fractional p -Laplacian would have the UCP, then similar methods as in [KRZ22] (see also [GKS16]) could be invoked to prove global uniqueness of the coefficients.

Conventions. Throughout this article, we denote by $B_r(x_0)$, $Q_r(x_0)$ for $r > 0$, $x_0 \in \mathbb{R}^n$ the open ball of radius r with center x_0 and the open cube of side length $2r$ with center x_0 . Moreover, if $x_0 = 0$, we simply write $B_r = B_r(0)$ and $Q_r = Q_r(0)$.

3. THE FORWARD PROBLEM AND DN MAP

In this section we first state an auxiliary lemma which will be of constant use in this work and then establish the well-posedness of the Dirichlet problem related to the fractional p -Laplace equations (1.1). Finally, we introduce the DN map and show that it induces a continuous map from the trace space to its dual.

3.1. Auxiliary lemma.

Lemma 3.1 (cf. [Sim78, eq. (2.2)], [GM75, Lemma 5.1-5.2] and [SZ12, Appendix A]). *Let $n \in \mathbb{N}$, $1 < p < \infty$, then there exists a constant $c_p > 0$ such that for all $x, y \in \mathbb{R}^n$, there holds*

$$(3.1) \quad (|x|^{p-2}x - |y|^{p-2}y) \cdot (x - y) \geq c_p|x - y|^p$$

if $p \geq 2$ and

$$(3.2) \quad (|x|^{p-2}x - |y|^{p-2}y) \cdot (x - y) \geq c_p \frac{|x - y|^2}{(|x| + |y|)^{2-p}}$$

if $1 < p < 2$. Moreover, for all $1 < p < \infty$, we have

$$(3.3) \quad ||\xi|^{p-2}\xi - |\eta|^{p-2}\eta| \leq C_p(|\xi| + |\eta|)^{p-2}|\xi - \eta|$$

for all $\xi, \eta \in \mathbb{R}^n$ and some constant $C_p > 0$.

3.2. Well-posedness. Next we prove well-posedness of the forward problem. In this work, we use the following notion of weak solutions:

Definition 3.2 (Weak solutions). *Let $1 < p < \infty$, $0 < s < 1$, $\Omega \subset \mathbb{R}^n$ be a bounded open set, and assume that $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the uniform ellipticity condition (1.2). For any $f \in W^{s,p}(\mathbb{R}^n)$, we say that $u \in W^{s,p}(\mathbb{R}^n)$ is a weak solution to*

$$(3.4) \quad \begin{cases} \operatorname{Div}_s(\sigma|d_s u|^{p-2}d_s u) = 0 & \text{in } \Omega, \\ u = f & \text{in } \Omega_e, \end{cases}$$

if u is a distributional solution and $u - f \in \widetilde{W}^{s,p}(\Omega)$.

Remark 3.3. Observe that if u is a distributional solution of (3.4), then there holds

$$\int_{\mathbb{R}^{2n}} \sigma(x, y) |u(x) - u(y)|^{p-2} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+sp}} dx dy = 0$$

for all $\varphi \in \widetilde{W}^{s,p}(\Omega)$.

Theorem 3.4 (Well-posedness). *Let $1 < p < \infty$, $0 < s < 1$, $\Omega \subset \mathbb{R}^n$ be a bounded open set, and assume that $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the uniform ellipticity condition (1.2). Then for any $f \in W^{s,p}(\mathbb{R}^n)$, there is a unique solution $u \in W^{s,p}(\mathbb{R}^n)$ of (3.4). In fact, it can be characterized as the unique minimizer of the fractional p -Dirichlet energy*

$$E_{s,p,\sigma}(v) := \int_{\mathbb{R}^{2n}} \sigma |d^s v|^p \frac{dx dy}{|x - y|^n}$$

over the class of all $v \in W^{s,p}(\mathbb{R}^n)$ with prescribed exterior data $v = f$ in Ω_e . Moreover, the unique solution satisfies the following estimate

$$(3.5) \quad [u]_{W^{s,p}(\mathbb{R}^n)} \leq C[f]_{W^{s,p}(\mathbb{R}^n)}$$

for some $C > 0$ depending only on λ and p .

Remark 3.5. Let us observe that we can construct the solution u as the unique minimizer of the functional $E_{s,p,\sigma}$ by the fact that our exterior condition f is such that

$$\int_{\Omega^c \times \Omega^c} \sigma(x, y) \frac{|f(x) - f(y)|^p}{|x - y|^{n+sp}} dx dy < \infty.$$

If the exterior condition would be less regular one should only integrate over $\mathbb{R}^{2n} \setminus (\Omega^c \times \Omega^c)$ and impose the exterior value in the sense that $u = f$ a.e. in Ω^c (cf. [RO16, Section 3]).

Before giving the proof of this well-posedness result, let us make the following elementary observation. The linear map $T: W^{s,p}(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n) \times L^p(\mathbb{R}^{2n})$ given by

$$u \mapsto \left(u, \frac{|u(x) - u(y)|}{|x - y|^{n/p+s}} \right),$$

where the target space is endowed with the usual product norm, is an isometry. Thus, by arguing as in [Bre11, Proposition 8.1], one sees that $W^{s,p}(\mathbb{R}^n)$ is separable in the range $1 \leq p < \infty$ and reflexive when $1 < p < \infty$. Now since $\widetilde{W}^{s,p}(\Omega)$ is a closed linear subspace of $W^{s,p}(\mathbb{R}^n)$ it follows that it has the same properties on the respective ranges.

Proof of Theorem 3.4. We proceed similarly as in [KRZ22, Theorem 5.8]. First we define the convex set

$$\widetilde{W}_f^{s,p}(\Omega) := \left\{ u \in W^{s,p}(\mathbb{R}^n) : u - f \in \widetilde{W}^{s,p}(\Omega) \right\} \subset W^{s,p}(\mathbb{R}^n),$$

and observe that it is weakly closed in the reflexive Banach space $W^{s,p}(\mathbb{R}^n)$. To see this, assume that $(u_k)_{k \in \mathbb{N}} \subset \widetilde{W}_f^{s,p}(\Omega)$ converges weakly to some $u \in W^{s,p}(\mathbb{R}^n)$ as $k \rightarrow \infty$. This implies that the sequence $(u_k - f)_{k \in \mathbb{N}} \subset \widetilde{W}^{s,p}(\Omega)$ converges weakly to $u - f \in W^{s,p}(\mathbb{R}^n)$.

Next, since weak limits are contained in the weak closure, the weak closure of convex sets coincide with the strong closure and $\widetilde{W}^{s,p}(\Omega)$ is by definition a closed subspace of $W^{s,p}(\mathbb{R}^n)$, we obtain that $u - f \in \widetilde{W}^{s,p}(\Omega)$. Hence, $\widetilde{W}_f^{s,p}(\Omega)$ is weakly closed in $W^{s,p}(\mathbb{R}^n)$. Next note that there holds

$$|a \pm b|^p \geq 2^{1-p} |a|^p - |b|^p$$

for all $a, b \in \mathbb{R}$. Hence, the uniform elliptic condition (1.2) of σ and the Poincaré inequality (Proposition 2.1) yield that

$$\begin{aligned} |E_{s,p,\sigma}(u)| &\geq \lambda \left(2^{1-p} [u - f]_{W^{s,p}(\mathbb{R}^n)}^p - [f]_{W^{s,p}(\mathbb{R}^n)}^p \right) \\ &\geq \frac{\lambda}{2^p} \left([u - f]_{W^{s,p}(\mathbb{R}^n)}^p + C \|u - f\|_{L^p(\mathbb{R}^n)}^p \right) - \lambda [f]_{W^{s,p}(\mathbb{R}^n)}^p \\ &\geq C \|u - f\|_{W^{s,p}(\mathbb{R}^n)}^p - \lambda [f]_{W^{s,p}(\mathbb{R}^n)}^p \\ &\geq C \|u\|_{W^{s,p}(\mathbb{R}^n)}^p - c \|f\|_{W^{s,p}(\mathbb{R}^n)}^p \end{aligned}$$

for all $u \in \widetilde{W}_f^{s,p}(\Omega)$ and some constants $C, c > 0$ only depending on λ and p .

Therefore, the functional $E_{s,p,\sigma}$ is coercive on $\widetilde{W}_f^{s,p}(\Omega)$, in the sense that

$$|E_{s,p,\sigma}(u)| \rightarrow \infty, \text{ when } \|u\|_{W^{s,p}(\mathbb{R}^n)} \rightarrow \infty,$$

for $u \in \widetilde{W}_f^{s,p}(\Omega)$. We also observe that by the assumptions on $\sigma(x, y)$, the functional $E_{s,p,\sigma}$ is convex and continuous on the closed, convex set $\widetilde{W}_f^{s,p}(\Omega) \subset W^{s,p}(\mathbb{R}^n)$. It is known that this implies that $E_{s,p,\sigma}$ is weakly lower semi-continuous on $\widetilde{W}_f^{s,p}(\Omega)$ (for example, see [BP12, Proposition 2.10]). Hence, using [Str08, Theorem 1.2] we see that there exists a minimizer $u \in \widetilde{W}_f^{s,p}(\Omega)$ of $E_{s,p,\sigma}$.

To proceed, let us show that the minimizer $u \in W^{s,p}(\mathbb{R}^n)$ solves (3.4) in the sense of distributions. Let $\varphi \in C_c^\infty(\Omega)$ then there holds $u_\varepsilon := u + \varepsilon\varphi \in \widetilde{W}_f^{s,p}(\Omega)$ for any $\varepsilon \in \mathbb{R}$. Moreover, by Hölder's inequality and the dominated convergence theorem, one can see that $E_{s,p,\sigma}$ is a C^1 -functional. Hence, the fact that u is a minimizer implies that there holds

$$\begin{aligned} 0 &= \left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} E_{s,p,\sigma}(u_\varepsilon) \\ &= p \int_{\mathbb{R}^{2n}} \sigma(x, y) |u(x) - u(y)|^{p-2} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|x - y|^{n+sp}} dx dy, \end{aligned}$$

and the claim follows. It remains to prove that the minimizer u is unique. Let us first show the assertion for the range $2 \leq p < \infty$ and then for $1 < p < 2$.

(i) First suppose that $2 \leq p < \infty$. Let $u, v \in W^{s,p}(\mathbb{R}^n)$ and set

$$\delta_{x,y}w := w(x) - w(y),$$

for any function $w: \mathbb{R}^n \rightarrow \mathbb{R}$. Using the estimate (3.1) of Lemma 3.1, we have the following strong monotonicity property

$$\begin{aligned} &\int_{\mathbb{R}^{2n}} \sigma (|\delta_{x,y}u|^{p-2} \delta_{x,y}u - |\delta_{x,y}v|^{p-2} \delta_{x,y}v) (\delta_{x,y}u - \delta_{x,y}v) \frac{dx dy}{|x - y|^{n+sp}} \\ (3.6) \quad &\geq \lambda c_p \int_{\mathbb{R}^{2n}} \frac{|\delta_{x,y}u - \delta_{x,y}v|^p}{|x - y|^{n+sp}} dx dy \\ &= \lambda c_p \int_{\mathbb{R}^{2n}} \frac{|(u(x) - u(y)) - (v(x) - v(y))|^p}{|x - y|^{n+sp}} dx dy \\ &= \lambda c_p [u - v]_{W^{s,p}(\mathbb{R}^n)}^p. \end{aligned}$$

Now, if $u, v \in W^{s,p}(\mathbb{R}^n)$ are solutions to (3.4) then $u - v \in \widetilde{W}_f^{s,p}(\Omega)$. Hence the left hand side vanishes and by the Poincaré inequality (Theorem 2.1) we can lower bound the right hand side by some positive multiple of $\|u - v\|_{L^p(\mathbb{R}^n)}^p$ but this gives $u = v$ a.e. in \mathbb{R}^n .

- (ii) Next let $1 < p < 2$. Using the same notation as before, we obtain by raising the estimate (3.2) of Lemma 3.1 to the power $p/2$ the bound

$$c_p^{p/2} |a - b|^p \leq [(|a|^{p-2}a - |b|^{p-2}b) \cdot (a - b)]^{p/2} (|a| + |b|)^{(2-p)\frac{p}{2}}$$

for all $a, b \in \mathbb{R}^n$. Now using Hölder's inequality with $\frac{2-p}{2} + \frac{p}{2} = 1$ and the uniform ellipticity of σ , we deduce

$$\begin{aligned}
& \lambda^{p/2} c_p^{p/2} [u - v]_{W^{s,p}(\mathbb{R}^n)}^p \\
& \leq \int_{\mathbb{R}^{2n}} |\delta_{x,y}u - \delta_{x,y}v|^p \frac{dxdy}{|x - y|^{n+sp}} \\
& \leq \lambda^{p/2} c_p^{p/2} \int_{\mathbb{R}^{2n}} [(|\delta_{x,y}u|^{p-2}\delta_{x,y}u - |\delta_{x,y}v|^{p-2}\delta_{x,y}v)(\delta_{x,y}u - \delta_{x,y}v)]^{p/2} \\
& \quad \cdot (|\delta_{x,y}u| + |\delta_{x,y}v|)^{(2-p)\frac{p}{2}} \frac{dxdy}{|x - y|^{n+sp}} \\
& \leq \lambda^{p/2} c_p^{p/2} \left\| (|\delta_{x,y}u| + |\delta_{x,y}v|)^{(2-p)\frac{p}{2}} \right\|_{L^{\frac{2}{2-p}}(\mathbb{R}^n; |x-y|^{-(n+sp)})} \\
(3.7) \quad & \cdot \left\| [(|\delta_{x,y}u|^{p-2}\delta_{x,y}u \right. \\
& \quad \left. - |\delta_{x,y}v|^{p-2}\delta_{x,y}v) (\delta_{x,y}u - \delta_{x,y}v)]^{p/2} \right\|_{L^{2/p}(\mathbb{R}^n; |x-y|^{-(n+sp)})} \\
& \leq c_p^{p/2} 2^{p-1} \left([u]_{W^{s,p}(\mathbb{R}^n)}^p + [v]_{W^{s,p}(\mathbb{R}^n)}^p \right)^{\frac{2-p}{2}} \\
& \quad \cdot \left(\int_{\mathbb{R}^{2n}} \sigma(|\delta_{x,y}u|^{p-2}\delta_{x,y}u \right. \\
& \quad \left. - |\delta_{x,y}v|^{p-2}\delta_{x,y}v) (\delta_{x,y}u - \delta_{x,y}v) \frac{dxdy}{|x - y|^{n+sp}} \right)^{p/2}.
\end{aligned}$$

Now, arguing as for the previous case $p \geq 2$, the right hand side vanishes if $u, v \in W^{s,p}(\mathbb{R}^n)$ are solutions to (3.4) as $u - v \in \widetilde{W}^{s,p}(\Omega)$ and the left hand side can be lower bounded by a positive multiples of $\|u - v\|_{L^p(\mathbb{R}^n)}^p$ by using the Poincaré inequality again. Hence, we can conclude that $u = v$ a.e. in \mathbb{R}^n .

To complete the proof, let us establish the estimate (3.5). By Remark 3.5, we can test the equation (3.4) with any $\varphi \in \widetilde{W}^{s,p}(\Omega)$ and in particular with $u - f$. Using the uniform ellipticity (1.2), writing $u = (u - f) + f$ and applying Young's inequality, we obtain

$$\begin{aligned}
\lambda [u]_{W^{s,p}(\mathbb{R}^n)}^p & \leq \int_{\mathbb{R}^{2n}} \sigma(x, y) \frac{|u(x) - u(y)|^p}{|x - y|^{n+sp}} dxdy \\
& \leq \int_{\mathbb{R}^{2n}} \sigma(x, y) |u(x) - u(y)|^{p-2} \frac{(u(x) - u(y))(u(x) - u(y))}{|x - y|^{n+sp}} dxdy \\
& = \int_{\mathbb{R}^{2n}} \sigma(x, y) |u(x) - u(y)|^{p-2} \frac{(u(x) - u(y))(f(x) - f(y))}{|x - y|^{n+sp}} dxdy \\
& \leq \lambda^{-1} \int_{\mathbb{R}^{2n}} \frac{|u(x) - u(y)|^{p-1} |f(x) - f(y)|}{|x - y|^{n+sp}} dxdy \\
& \leq \underbrace{\epsilon [u]_{W^{s,p}(\mathbb{R}^n)}^p + C_\epsilon \lambda^{-p} [f]_{W^{s,p}(\mathbb{R}^n)}^p}_{ab \leq \epsilon a^p + C_\epsilon b^{p'} \text{ with } 1/p + 1/p' = 1}
\end{aligned}$$

for any $\epsilon > 0$, where $C_\epsilon > 0$ is a constant depending on ϵ . In particular, by choosing $\epsilon = \lambda/2$, we obtain $[u]_{W^{s,p}(\mathbb{R}^n)} \leq C[f]_{W^{s,p}(\mathbb{R}^n)}$ for some $C > 0$ depending only on λ and p . This proves the assertion. \square

Next we introduce the abstract trace space:

Definition 3.6. *Let $\Omega \subset \mathbb{R}^n$ be an open set, $0 < s < 1$ and $1 < p < \infty$. Then the abstract trace space $X^{s,p}(\Omega)$ is given by $X^{s,p}(\Omega) := W^{s,p}(\mathbb{R}^n)/\widetilde{W}^{s,p}(\Omega)$, and we endow it with the quotient norm*

$$\|f\|_{X^{s,p}(\Omega)} := \inf_{u \in \widetilde{W}^{s,p}(\Omega)} \|f - u\|_{W^{s,p}(\mathbb{R}^n)}.$$

Remark 3.7. *Let us point out:*

- (i) *In the above definition and later on we simply write f for an element in $X^{s,p}(\Omega)$ instead of the more precise notation $[f]$. Note that since $\widetilde{W}^{s,p}(\Omega)$ is a closed subspace of the Banach space $W^{s,p}(\mathbb{R}^n)$, the abstract trace space is again a separable, reflexive Banach space in the respective ranges $1 \leq p < \infty$ and $1 < p < \infty$.*
- (ii) *If $\Omega \subset \mathbb{R}^n$ has a bounded Lipschitz continuous boundary then any $f \in W^{s,p}(\Omega_e)$ with $\text{dist}(\text{supp } f, \partial\Omega) > 0$ corresponds to a unique equivalence class in $X^{s,p}(\Omega)$ and the quotient norm is equivalent to the $\|\cdot\|_{W^{s,p}(\Omega_e)}$ norm. In fact, by [CRTZ22, Lemma 3.2] the zero extension \bar{f} of f belongs to $W^{s,p}(\mathbb{R}^n)$ and satisfies $\|\bar{f}\|_{W^{s,p}(\mathbb{R}^n)} \leq C\|f\|_{W^{s,p}(\Omega_e)}$. Hence, we implicitly identify below f with the equivalence class $[\bar{f}]$. Moreover, by definition we have $\|\bar{f}\|_{X^{s,p}(\Omega)} \leq \|\bar{f}\|_{W^{s,p}(\mathbb{R}^n)} \leq C\|f\|_{W^{s,p}(\Omega_e)}$. On the other hand, there exists a sequence $u_k \in \widetilde{W}^{s,p}(\Omega)$, $k \in \mathbb{N}$, such that $\|f - u_k\|_{W^{s,p}(\mathbb{R}^n)} \rightarrow \|\bar{f}\|_{X^{s,p}(\Omega)}$ as $k \rightarrow \infty$. Since $\partial\Omega$ has measure zero, we know that u_k vanish a.e. in Ω_e and thus there holds*

$$\|f\|_{W^{s,p}(\Omega_e)} = \|f - u_k\|_{W^{s,p}(\Omega_e)} \leq \|\bar{f} - u_k\|_{W^{s,p}(\mathbb{R}^n)} \rightarrow \|\bar{f}\|_{X^{s,p}(\Omega)}$$

as $k \rightarrow \infty$. This shows that these two norms are equivalent.

We have the following uniqueness result:

Corollary 3.8. *Let $1 < p < \infty$, $0 < s < 1$, $\Omega \subset \mathbb{R}^n$ a bounded open set and assume that $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the uniform ellipticity condition (1.2). Let $u_j \in W^{s,p}(\mathbb{R}^n)$ be the unique solutions of (3.4) with exterior values $f_j \in W^{s,p}(\mathbb{R}^n)$ for $j = 1, 2$. If $f_1 - f_2 \in \widetilde{W}^{s,p}(\Omega)$, then $u_1 = u_2$.*

Proof. By assumption we have

$$u_1 - u_2 = (u_1 - f_1) - (u_2 - f_2) + (f_1 - f_2) \in \widetilde{W}^{s,p}(\Omega).$$

Hence, arguing as in the proof of Theorem 3.4, the strong monotonicity properties (3.6) for $p \geq 2$, and (3.7) for $1 < p < 2$, respectively, show that $u_1 = u_2$. \square

3.3. DN maps. With Theorem 3.4 at hand, we can introduce the DN map Λ_σ to formulate the inverse problem. If $f \in W^{s,p}(\mathbb{R}^n)$, then the DN map is formally defined by

$$(3.8) \quad \Lambda_\sigma(f) = \text{Div}_s(\sigma|d_s u_f|^{p-2} d_s u_f)|_{\Omega_e},$$

where $u_f \in W^{s,p}(\mathbb{R}^n)$ is the unique solution of

$$(3.9) \quad \begin{cases} \text{Div}_s(\sigma|d_s u|^{p-2} d_s u) = 0 & \text{in } \Omega, \\ u = f & \text{in } \Omega_e, \end{cases}$$

(cf. Theorem 3.4). As the solution u_f is usually not regular enough to justify the pointwise definition (3.8), we define it in general in the distributional sense by

$$(3.10) \quad \langle \Lambda_\sigma(f), g \rangle := \int_{\Omega_e} \Lambda_\sigma(f) g \, dx := \int_{\mathbb{R}^{2n}} \sigma |d_s u|^{p-2} d_s u d_s g \frac{dxdy}{|x-y|^n},$$

where $f, g \in X^{s,p}(\Omega)$. We have:

Proposition 3.9 (DN maps). *Let $1 < p < \infty$, $0 < s < 1$, $\Omega \subset \mathbb{R}^n$ a bounded open set and assume that $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the uniform ellipticity condition (1.2). Then the DN map Λ_σ introduced via (3.10) is a well-defined operator from $X^{s,p}(\Omega)$ to $(X^{s,p}(\Omega))^*$ and satisfies the estimate*

$$\|\Lambda_\sigma(f)\|_{(X^{s,p}(\Omega))^*} \leq C \|f\|_{X^{s,p}(\Omega)}^{p-1}$$

for all $f \in X^{s,p}(\Omega)$ and some $C > 0$. Here $(X^{s,p}(\Omega))^*$ denotes the dual space of $X^{s,p}(\Omega)$.

Proof. First note that by Corollary 3.8 for any $f \in X^{s,p}(\Omega)$, there is a unique solution $u_f \in W^{s,p}(\mathbb{R}^n)$ of (3.9). Moreover, changing in the weak formulation of the DN map (3.10), the function $g \in W^{s,p}(\mathbb{R}^n)$ to $g + \varphi$ with $\varphi \in \widetilde{W}^{s,p}(\Omega)$ does not change the value of the DN map as by construction u_f solves (3.9). Hence, the DN map Λ_σ is well-defined. Finally, by the Hölder's inequality with $\frac{p-1}{p} + \frac{1}{p} = 1$, we have

$$\begin{aligned} |\langle \Lambda_\sigma(f), g \rangle| &\leq C \|d_s u\|_{L^p(\mathbb{R}^{2n}, |x-y|^{-n})}^{p-1} \|d_s g\|_{L^p(\mathbb{R}^{2n}, |x-y|^{-n})} \\ &\leq C \|f\|_{W^{s,p}(\mathbb{R}^n)}^{p-1} \|g\|_{W^{s,p}(\mathbb{R}^n)} \end{aligned}$$

for all $f, g \in W^{s,p}(\mathbb{R}^n)$, for some constant $C > 0$ independent of f and g . Hence, taking the infimum over all representations of $f, g \in X^{s,p}(\Omega)$ and dividing by $\|g\|_{X^{s,p}(\Omega)}$, we obtain

$$\|\Lambda_\sigma(f)\|_{(X^{s,p}(\Omega))^*} \leq C \|f\|_{X^{s,p}(\Omega)}^{p-1}.$$

This completes the proof. \square

4. EXTERIOR RECONSTRUCTION

In this section, we establish the exterior reconstruction result, which is the main theorem of this article.

Lemma 4.1 (Exterior conditions). *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $0 < s < 1$, $1 \leq p < \infty$ and $x_0 \in W \subset \Omega_e$ for an open set W . There exists a sequence $(\Phi_N)_{N \in \mathbb{N}} \subset C_c^\infty(W)$ such that*

- (i) for all $N \in \mathbb{N}$ it holds $[\Phi_N]_{W^{s,p}(\mathbb{R}^n)} = 1$,
- (ii) for all $0 \leq t < s$ there holds $\|\Phi_N\|_{W^{t,p}(\mathbb{R}^n)} \rightarrow 0$ as $N \rightarrow \infty$
- (iii) and $\text{supp}(\Phi_N) \rightarrow \{x_0\}$ as $N \rightarrow \infty$.

Remark 4.2. *Kohn and Vogelius proved a similar result in their celebrated work on boundary determination for the conductivity equation (cf. [KV84, Lemma 1]) for the Sobolev spaces $H^s(\partial\Omega)$, where $\Omega \subset \mathbb{R}^n$ is an open bounded set with smooth boundary (see also [CRZ22, Lemma 5.5]).*

Proof of Lemma 4.1. By translation and scaling, we may assume that $Q_1 \subset W$ and $x_0 = 0$ without loss of generality. Similarly, as in [KV84], we choose any nonzero $\psi \in C_c^\infty(\mathbb{R})$ with $\text{supp}(\psi) \subset (-1, 1)$ and let Ψ be the n -fold tensor product of ψ , that is $\Psi(x) = \prod_{k=1}^n \psi(x_k)$ with $x = (x_1, x_2, \dots, x_n)$. Next we define the

sequence $(\Psi_N)_{N \in \mathbb{N}}$ by $\Psi_N(x) := \Psi(Nx)$. We clearly have $\Psi_N \in C_c^\infty(Q_{1/N})$ and $\text{supp}(\Psi_N) \rightarrow \{0\}$. Moreover, by a simple change of variables we have

$$(4.1) \quad \|\Psi_N\|_{L^p(\mathbb{R}^n)} = N^{-n/p} \|\Psi\|_{L^p(\mathbb{R}^n)}$$

and

$$(4.2) \quad [\Psi_N]_{W^{t,p}(\mathbb{R}^n)} = N^{t-n/p} [\Psi]_{W^{t,p}(\mathbb{R}^n)}$$

for all $0 < t < 1$, $1 \leq p < \infty$ and $N \in \mathbb{N}$. Observe that $[\Psi]_{W^{t,p}(\mathbb{R}^n)} > 0$ for all $0 \leq t < 1$. This is an immediate consequence of $0 \neq \psi \in C_c^\infty((-1, 1))$ and the Poincaré inequality (Theorem 2.1). Thus, for all $0 \leq t < 1$, $1 \leq p < \infty$ there exist constants $C_{t,p}, C'_{t,p} > 0$ such that

$$C'_{t,p} N^{t-n/p} \leq \|\Psi_N\|_{W^{t,p}(\mathbb{R}^n)} \leq C_{t,p} N^{t-n/p}$$

for all $N \in \mathbb{N}$.

Finally, we introduce for $N \in \mathbb{N}$ the rescaled functions Φ_N by

$$\Phi_N := \frac{\Psi_N}{[\Psi_N]_{W^{s,p}(\mathbb{R}^n)}} \in C_c^\infty(Q_{1/N}).$$

We clearly have $[\Phi_N]_{W^{s,p}(\mathbb{R}^n)} = 1$ and $\text{supp}(\Phi_N) \rightarrow \{0\}$. Moreover, from (4.1) and (4.2) we deduce that

$$\|\Phi_N\|_{L^p(\mathbb{R}^n)} = \frac{\|\Psi_N\|_{L^p(\mathbb{R}^n)}}{[\Psi_N]_{W^{s,p}(\mathbb{R}^n)}} = \frac{N^{-n/p} \|\Psi\|_{L^p(\mathbb{R}^n)}}{N^{s-n/p} [\Psi]_{W^{s,p}(\mathbb{R}^n)}} = N^{-s} \frac{\|\Psi\|_{L^p(\mathbb{R}^n)}}{[\Psi]_{W^{s,p}(\mathbb{R}^n)}} \rightarrow 0$$

and

$$[\Phi_N]_{W^{t,p}(\mathbb{R}^n)} = \frac{[\Psi_N]_{W^{t,p}(\mathbb{R}^n)}}{[\Psi_N]_{W^{s,p}(\mathbb{R}^n)}} = \frac{N^{t-n/p} [\Psi]_{W^{t,p}(\mathbb{R}^n)}}{N^{s-n/p} [\Psi]_{W^{s,p}(\mathbb{R}^n)}} = N^{t-s} \frac{[\Psi]_{W^{t,p}(\mathbb{R}^n)}}{[\Psi]_{W^{s,p}(\mathbb{R}^n)}} \rightarrow 0$$

as $N \rightarrow \infty$, when $0 < t < s$. Hence, the sequence $(\Phi_N)_{N \in \mathbb{N}}$ satisfies the properties (i) – (iii). \square

Lemma 4.3. (Energy concentration property) *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $0 < s < 1$, $1 < p < \infty$ and $x_0 \in W \Subset \Omega_e$. Assume that $(\Phi_N)_{N \in \mathbb{N}} \subset C_c^\infty(W)$ is a sequence satisfying the properties (i) – (iii) of Lemma 4.1. If $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfies the uniform ellipticity condition (1.2), $(\sigma(x, \cdot): W \rightarrow \mathbb{R})_{x \in \mathbb{R}^n}$ is equicontinuous at x_0 and $\sigma(\cdot, x_0) \in C(W)$, then we have*

$$\sigma(x_0, x_0) = \lim_{N \rightarrow \infty} E_{s,p,\sigma}(\Phi_N).$$

Remark 4.4. *We remark that if $\sigma(x, y) = \alpha(x)\beta(y)$ for some uniformly elliptic functions $\alpha, \beta \in L^\infty(\mathbb{R}^n) \cap C(W)$ then σ satisfies the assumptions of Lemma 4.3.*

Proof of Lemma 4.3. First note that we can decompose $\sigma(x, y)$ as

$$\sigma(x, y) = (\sigma(x, y) - \sigma(x, x_0)) + (\sigma(x, x_0) - \sigma(x_0, x_0)) + \sigma(x_0, x_0), \text{ for all } x, y \in \mathbb{R}^n.$$

This implies

$$(4.3) \quad \begin{aligned} E_{s,p,\sigma}(\Phi_N) &= \int_{\mathbb{R}^{2n}} (\sigma(x, y) - \sigma(x, x_0)) |d_s \Phi_N|^p \frac{dxdy}{|x-y|^n} \\ &\quad + \int_{\mathbb{R}^{2n}} (\sigma(x, x_0) - \sigma(x_0, x_0)) |d_s \Phi_N|^p \frac{dxdy}{|x-y|^n} \\ &\quad + \sigma(x_0, x_0) \int_{\mathbb{R}^{2n}} |d_s \Phi_N|^p \frac{dxdy}{|x-y|^n}. \end{aligned}$$

By (i) of Lemma 4.1, it follows that the last term is equal to $\sigma(x_0, x_0)$. Hence, to establish the assertion it suffice to prove that the two remaining integrals go to zero as $N \rightarrow \infty$.

Next observe that by the product rule for the fractional gradient (2.2),

$$|d_s(\Phi_N \psi)|^p \leq C (|\Phi_N(x)|^p |d_s \psi(x, y)|^p + |\psi(y)|^p |d_s \Phi_N(x, y)|^p).$$

If $\psi \in C_b^1(\mathbb{R}^n)$, then the mean value theorem yields that

$$\begin{aligned} & \int_{\mathbb{R}^n} |d_s \psi(x, y)|^p \frac{dy}{|x-y|^n} \\ & \lesssim \int_{B_1(x)} \frac{|\psi(x) - \psi(y)|^p}{|x-y|^{n+sp}} dy + \int_{\mathbb{R}^n \setminus B_1(x)} \frac{|\psi(x) - \psi(y)|^p}{|x-y|^{n+sp}} dy \\ & \lesssim \|\nabla \psi\|_{L^\infty(\mathbb{R}^n)}^p \int_{B_1(x)} \frac{dy}{|x-y|^{n+p(s-1)}} + \|\psi\|_{L^\infty(\mathbb{R}^n)}^p \int_{\mathbb{R}^n \setminus B_1(x)} \frac{dy}{|x-y|^{n+sp}} \\ & \lesssim \left(\int_{B_1(0)} \frac{dz}{|z|^{n+p(s-1)}} + \int_{\mathbb{R}^n \setminus B_1(0)} \frac{dz}{|z|^{n+ps}} \right) \|\psi\|_{C^1(\mathbb{R}^n)}^p \\ & \lesssim \|\psi\|_{C^1(\mathbb{R}^n)}^p, \end{aligned}$$

for all $x \in \mathbb{R}^n$. By (ii) of Lemma 4.1, we have $\|\Phi_N\|_{L^p(\mathbb{R}^n)} \rightarrow 0$ as $N \rightarrow \infty$, and thus for all $\psi \in C_b^1(\mathbb{R}^n)$ there holds

$$(4.4) \quad \int_{\mathbb{R}^{2n}} |\Phi_N(x)|^p |d_s \psi(x, y)|^p \frac{dx dy}{|x-y|^n} \lesssim \|\psi\|_{C^1(\mathbb{R}^n)}^p \|\Phi_N\|_{L^p(\mathbb{R}^n)}^p \rightarrow 0$$

as $N \rightarrow \infty$. Consider now a sequence of functions $(\eta_M)_{M \in \mathbb{N}} \subset C_c^1(\mathbb{R}^n)$ such that for all $M \in \mathbb{N}$ it holds $0 \leq \eta_M \leq 1$, $\eta_M|_{Q_{1/2M}(x_0)} = 1$ and $\eta_M|_{(Q_{1/M}(x_0))^c} = 0$. If $N \in \mathbb{N}$ is sufficiently large, then $\eta_M \Phi_N = \Phi_N$ and hence by using (4.4), we deduce

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \left| \int_{\mathbb{R}^{2n}} (\sigma(x, y) - \sigma(x, x_0)) |d_s \Phi_N|^p \frac{dx dy}{|x-y|^n} \right| \\ & = \limsup_{N \rightarrow \infty} \left| \int_{\mathbb{R}^{2n}} (\sigma(x, y) - \sigma(x, x_0)) |d_s(\eta_M \Phi_N)|^p \frac{dx dy}{|x-y|^n} \right| \\ & \lesssim \limsup_{N \rightarrow \infty} \left| \int_{\mathbb{R}^{2n}} (\sigma(x, y) - \sigma(x, x_0)) |\eta_M(y)|^p |d_s \Phi_N|^p \frac{dx dy}{|x-y|^n} \right| \\ & \lesssim \sup_{y \in Q_{1/M}(x_0)} \sup_{x \in \mathbb{R}^n} |\sigma(x, y) - \sigma(x, x_0)| \end{aligned}$$

for all $M \in \mathbb{N}$, where we used (i) of Lemma 4.1 in the last step. Using the equicontinuity assumption on σ , we deduce that

$$\limsup_{N \rightarrow \infty} \left| \int_{\mathbb{R}^{2n}} (\sigma(x, y) - \sigma(x, x_0)) |d_s \Phi_N|^p \frac{dx dy}{|x-y|^n} \right| = 0.$$

Similarly, we have

$$\begin{aligned} & \limsup_{N \rightarrow \infty} \left| \int_{\mathbb{R}^{2n}} (\sigma(x, x_0) - \sigma(x_0, x_0)) |d_s \Phi_N|^p \frac{dx dy}{|x-y|^n} \right| \\ & = \limsup_{N \rightarrow \infty} \left| \int_{\mathbb{R}^{2n}} (\sigma(x, x_0) - \sigma(x_0, x_0)) |d_s(\eta_M \Phi_N)|^p \frac{dx dy}{|x-y|^n} \right| \\ & \lesssim \limsup_{N \rightarrow \infty} \left| \int_{\mathbb{R}^{2n}} (\sigma(x, x_0) - \sigma(x_0, x_0)) |\eta_M(x)|^p |d_s \Phi_N|^p \frac{dx dy}{|x-y|^n} \right| \\ & \lesssim \sup_{x \in Q_{1/M}(x_0)} |\sigma(x, x_0) - \sigma(x_0, x_0)| \end{aligned}$$

for all $M \in \mathbb{N}$. By the continuity of $x \mapsto \sigma(x, x_0)$ it follows that the last term vanishes as $M \rightarrow \infty$. Hence, taking the limit $N \rightarrow \infty$ in (4.3), we obtain

$$\sigma(x_0, x_0) = \lim_{N \rightarrow \infty} E_{s,p,\sigma}(\Phi_N).$$

This completes the proof. \square

Now, we observe that if u_N denotes the unique solution of

$$\begin{cases} \operatorname{Div}_s(\sigma|d_s u|^{p-2}d_s u) = 0 & \text{in } \Omega, \\ u = \Phi_N & \text{in } \Omega_e, \end{cases}$$

then by writing $u_N = (u_N - \Phi_N) + \Phi_N \in \widetilde{W}^{s,p}(\Omega) + W^{s,p}(\mathbb{R}^n)$ there holds

$$\int_{\mathbb{R}^{2n}} \sigma |d_s u_N|^p \frac{dxdy}{|x-y|^n} = \int_{\mathbb{R}^{2n}} \sigma |d_s u_N|^{p-2} d_s u_N d_s \Phi_N \frac{dxdy}{|x-y|^n}.$$

Thus, by Proposition 3.9 we obtain

$$\langle \Lambda_\sigma \Phi_N, \Phi_N \rangle = \int_{\mathbb{R}^{2n}} \sigma |d_s u_N|^p \frac{dxdy}{|x-y|^n} = \int_{\mathbb{R}^{2n}} \sigma |d_s u_N|^{p-2} d_s u_N d_s \Phi_N \frac{dxdy}{|x-y|^n},$$

so that

$$\begin{aligned} & \langle \Lambda_\sigma \Phi_N, \Phi_N \rangle \\ &= \int_{\mathbb{R}^{2n}} \sigma |d_s \Phi_N|^p \frac{dxdy}{|x-y|^n} \\ (4.5) \quad &+ \int_{\mathbb{R}^{2n}} \sigma \left(|d_s u_N|^{p-2} d_s u_N - |d_s \Phi_N|^{p-2} d_s \Phi_N \right) d_s \Phi_N \frac{dxdy}{|x-y|^n} \\ &= E_{s,p,\sigma}(\Phi_N) \\ &+ \int_{\mathbb{R}^{2n}} \sigma \left(|d_s u_N|^{p-2} d_s u_N - |d_s \Phi_N|^{p-2} d_s \Phi_N \right) d_s \Phi_N \frac{dxdy}{|x-y|^n}. \end{aligned}$$

We will recover the value of σ at (x_0, x_0) by showing that the second term goes to zero as $N \rightarrow \infty$. For this purpose, we show next:

Lemma 4.5. *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $0 < s < 1$, $1 < p < \infty$ and $x_0 \in W \Subset \Omega_e$. Assume that $(\Phi_N)_{N \in \mathbb{N}} \subset C_c^\infty(W)$ is a sequence satisfying the properties (i) – (iii) of Lemma 4.1. Let $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfy the uniform elliptic condition (1.2). Then we have*

$$(4.6) \quad \|u_N - \Phi_N\|_{W^{s,p}(\mathbb{R}^n)} \rightarrow 0$$

when $N \rightarrow \infty$, where $u_N \in W^{s,p}(\mathbb{R}^n)$ is the unique solution to

$$(4.7) \quad \begin{cases} \operatorname{Div}_s(\sigma|d_s u|^{p-2}d_s u) = 0 & \text{in } \Omega, \\ u = \Phi_N & \text{in } \Omega_e. \end{cases}$$

Proof. We divide the proof into the following two cases:

(i) For $2 \leq p < \infty$:

By the strong monotonicity property (3.6), there holds

$$\begin{aligned} [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)}^p &\leq C \int_{\mathbb{R}^{2n}} \sigma (|\delta_{x,y} u_N|^{p-2} \delta_{x,y} u_N \\ &\quad - |\delta_{x,y} \Phi_N|^{p-2} \delta_{x,y} \Phi_N) (\delta_{x,y} u_N - \delta_{x,y} \Phi_N) \frac{dxdy}{|x-y|^{n+sp}}, \end{aligned}$$

for all $N \in \mathbb{N}$ and some $C > 0$ only depending on n, p, λ . Next let us introduce the auxiliary function $\widetilde{\Phi}_N := u_N - \Phi_N \in \widetilde{W}^{s,p}(\Omega)$. Hence using that u_N solves (4.7) as well as that $\Phi_N \in C_c^\infty(W)$ and $\widetilde{\Phi}_N$ have disjoint

supports, we get

$$\begin{aligned}
& [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)}^p \\
& \leq C \int_{\mathbb{R}^{2n}} \sigma (|\delta_{x,y} u_N|^{p-2} \delta_{x,y} u_N - |\delta_{x,y} \Phi_N|^{p-2} \delta_{x,y} \Phi_N) \delta_{x,y} \tilde{\Phi}_N \frac{dxdy}{|x-y|^{n+sp}} \\
& = -C \underbrace{\int_{\mathbb{R}^{2n}} \sigma |\delta_{x,y} \Phi_N|^{p-2} \delta_{x,y} \Phi_N \delta_{x,y} \tilde{\Phi}_N \frac{dxdy}{|x-y|^{n+sp}}}_{\text{since } u_N \text{ solves (4.7)}} \\
& = -C \int_{\mathbb{R}^{2n}} \sigma |\delta_{x,y} \Phi_N|^{p-2} \left(\Phi_N(x) \tilde{\Phi}_N(x) + \Phi_N(y) \tilde{\Phi}_N(y) \right. \\
& \quad \left. - \Phi_N(x) \tilde{\Phi}_N(y) - \Phi_N(y) \tilde{\Phi}_N(x) \right) \frac{dxdy}{|x-y|^{n+sp}} \\
& = C \int_{\mathbb{R}^{2n}} \sigma |\delta_{x,y} \Phi_N|^{p-2} \left(\Phi_N(x) \tilde{\Phi}_N(y) + \Phi_N(y) \tilde{\Phi}_N(x) \right) \frac{dxdy}{|x-y|^{n+sp}} \\
& := C(I_1 + I_2),
\end{aligned}$$

where

$$\begin{aligned}
I_1 & := \int_{\mathbb{R}^{2n}} \sigma |\delta_{x,y} \Phi_N|^{p-2} \Phi_N(x) \tilde{\Phi}_N(y) \frac{dxdy}{|x-y|^{n+sp}}, \\
I_2 & := \int_{\mathbb{R}^{2n}} \sigma |\delta_{x,y} \Phi_N|^{p-2} \Phi_N(y) \tilde{\Phi}_N(x) \frac{dxdy}{|x-y|^{n+sp}}.
\end{aligned}$$

By Hölder's inequality, the convexity of $x \mapsto |x|^q$ for $q > 1$, and Minkowski's inequality, we obtain

$$\begin{aligned}
|I_1| & = \left| \int_{\mathbb{R}^{2n}} \sigma |\delta_{x,y} \Phi_N|^{p-2} \Phi_N(x) \tilde{\Phi}_N(y) \frac{dxdy}{|x-y|^{n+sp}} \right| \\
& \leq C \int_W \int_{\Omega} \sigma (|\Phi_N(x)|^{p-2} + |\Phi_N(y)|^{p-2}) |\Phi_N(x)| |\tilde{\Phi}_N(y)| \frac{dydx}{|x-y|^{n+sp}} \\
& = C \|\sigma\|_{L^\infty(W \times \Omega)} \int_W \int_{\Omega} |\Phi_N(x)|^{p-1} |\tilde{\Phi}_N(y)| \frac{dydx}{|x-y|^{n+sp}} \\
& \leq C \|\sigma\|_{L^\infty(W \times \Omega)} \int_W |\Phi_N(x)|^{p-1} \left(\int_{\Omega} \frac{|\tilde{\Phi}_N(y)|}{|x-y|^{n+sp}} dy \right) dx \\
& \leq C \|\sigma\|_{L^\infty(W \times \Omega)} \|\Phi_N\|_{L^{\frac{p}{p-1}}(W)}^{p-1} \left\| \int_{\Omega} \frac{|\tilde{\Phi}_N(y)|}{|x-y|^{n+sp}} dy \right\|_{L^p(W)} \\
& \leq C \|\sigma\|_{L^\infty(W \times \Omega)} \|\Phi_N\|_{L^p(W)}^{p-1} \int_{\Omega} |\tilde{\Phi}_N(y)| \left(\int_W \frac{dx}{|x-y|^{(n+sp)p}} \right)^{1/p} dy \\
& = C \|\sigma\|_{L^\infty(W \times \Omega)} \|\Phi_N\|_{L^p(W)}^{p-1} \int_{\Omega} |\tilde{\Phi}_N(y)| \left(\int_W \frac{dx}{|x-y|^{(n+sp)p}} \right)^{1/p} dy.
\end{aligned}$$

Now define $d := \text{dist}(\Omega, W) > 0$. Hence, Hölder's and Young's inequality imply

$$\begin{aligned}
|I_1| & \leq C \|\sigma\|_{L^\infty(W \times \Omega)} \|\Phi_N\|_{L^p(W)}^{p-1} |\Omega|^{\frac{p-1}{p}} \|\tilde{\Phi}_N\|_{L^p(\Omega)} \frac{|W|^{1/p}}{d^{n+sp}} \\
& \leq \varepsilon \|\tilde{\Phi}_N\|_{L^p(\Omega)}^p + C_\varepsilon \|\sigma\|_{L^\infty(W \times \Omega)}^{\frac{p}{p-1}} \|\Phi_N\|_{L^p(W)}^p |\Omega| \frac{|W|^{\frac{1}{p-1}}}{d^{\frac{p(n+sp)}{p-1}}},
\end{aligned}$$

for all $\varepsilon > 0$. The same estimate holds for I_2 after replacing $\|\sigma\|_{L^\infty(W \times \Omega)}$ by $\|\sigma\|_{L^\infty(\Omega \times W)}$. Hence, if we choose $\varepsilon > 0$ sufficiently small, then by using

Poincaré's inequality and recalling that $\tilde{\Phi}_N = u_N - \Phi_N$, the first term on the right hand side can be absorbed on the left hand side to obtain

$$\|u_N - \Phi_N\|_{W^{s,p}(\mathbb{R}^n)}^p \leq C_\varepsilon \|\sigma\|_{L^\infty((W \cup \Omega) \times (W \cup \Omega))}^{\frac{p}{p-1}} \|\Phi_N\|_{L^p(W)}^p |\Omega| \frac{|W|^{\frac{1}{p-1}}}{d^{\frac{p(n+sp)}{p-1}}}.$$

Now this expression goes to zero as N goes to ∞ by Lemma 4.1, (ii).

(ii) For $1 < p < 2$:

Applying this time the strong monotonicity property (3.7) on $u_N - \Phi_N$ gives

$$\begin{aligned} & [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)}^p \\ & \leq C ([u_N]_{W^{s,p}(\mathbb{R}^n)}^p + [\Phi_N]_{W^{s,p}(\mathbb{R}^n)}^p)^{\frac{2-p}{2}} \\ & \quad \cdot \left(\int_{\mathbb{R}^{2n}} \sigma (|\delta_{x,y} u_N|^{p-2} \delta_{x,y} u_N \right. \\ & \quad \left. - |\delta_{x,y} \Phi_N|^{p-2} \delta_{x,y} \Phi_N) (\delta_{x,y} u_N - \delta_{x,y} \Phi_N) \frac{dxdy}{|x-y|^{n+sp}} \right)^{p/2}, \end{aligned}$$

for some $C > 0$ only depending on n, p and λ . As in the previous case, this implies

$$\begin{aligned} & [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)}^p \\ & \leq C \|\sigma\|_{L^\infty((W \cup \Omega) \times (\Omega \cup W))} \left(\int_{\mathbb{R}^{2n}} |\delta_{x,y} \Phi_N|^{p-2} (|\Phi_N(x)| |\tilde{\Phi}_N(y)| \right. \\ & \quad \left. + |\Phi_N(y)| |\tilde{\Phi}_N(x)|) \frac{dxdy}{|x-y|^{n+sp}} \right)^{p/2} \\ & \leq C \|\sigma\|_{L^\infty((W \cup \Omega) \times (\Omega \cup W))} \left(\int_{\Omega} \int_W |\delta_{x,y} \Phi_N|^{p-2} |\Phi_N(x)| |\tilde{\Phi}_N(y)| \frac{dxdy}{|x-y|^{n+sp}} \right)^{p/2} \\ & = C \|\sigma\|_{L^\infty((W \cup \Omega) \times (\Omega \cup W))} \left(\int_{\Omega} \int_W |\Phi_N(x)|^{p-1} |\tilde{\Phi}_N(y)| \frac{dxdy}{|x-y|^{n+sp}} \right)^{p/2}, \end{aligned}$$

where we again set $\tilde{\Phi}_N = u_N - \Phi_N$. Here we used that u_N solves (4.7), σ is uniformly elliptic, $[\Phi_N]_{W^{s,p}(\mathbb{R}^n)} = 1$ and $\Phi_N, \tilde{\Phi}_N$ have disjoint supports. As in the previous case, this integral can be bounded from above as

$$\begin{aligned} [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)}^p & \leq C \|\sigma\|_{L^\infty((W \cup \Omega) \times (\Omega \cup W))} \\ & \quad \cdot \left(\|\Phi_N\|_{L^p(W)}^{p-1} |\Omega|^{\frac{p-1}{p}} \|\tilde{\Phi}_N\|_{L^p(\Omega)} \frac{|W|^{1/p}}{d^{n+sp}} \right)^{p/2}. \end{aligned}$$

Applying Young's inequality in the form $ab \leq \varepsilon a^2 + C_\varepsilon b^2$ gives

$$\begin{aligned} [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)}^p & \leq \varepsilon \|\tilde{\Phi}_N\|_{L^p(\Omega)}^p \\ & \quad + C_\varepsilon \|\sigma\|_{L^\infty((W \cup \Omega) \times (\Omega \cup W))}^2 \left(\|\Phi_N\|_{L^p(W)}^{p-1} |\Omega|^{\frac{p-1}{p}} \frac{|W|^{1/p}}{d^{n+sp}} \right)^p \end{aligned}$$

for all $\varepsilon > 0$. Using Poincaré's inequality and choosing ε sufficiently small, we can absorb the first term on the left hand side to obtain

$$\|u_N - \Phi_N\|_{W^{s,p}(\mathbb{R}^n)}^p \leq C_\varepsilon \|\sigma\|_{L^\infty((W \cup \Omega) \times (\Omega \cup W))}^2 \left(\|\Phi_N\|_{L^p(W)}^{p-1} |\Omega|^{\frac{p-1}{p}} \frac{|W|^{1/p}}{d^{n+sp}} \right)^p.$$

By Lemma 4.1, (ii) again, we deduce that $\|u_N - \Phi_N\|_{W^{s,p}(\mathbb{R}^n)} \rightarrow 0$ as $N \rightarrow \infty$.

This completes the proof. \square

Proposition 4.6. *Suppose that the assumptions of Lemma 4.5 hold, then we have*

$$\lim_{N \rightarrow \infty} \int_{\mathbb{R}^{2n}} \sigma \left(|d_s u_N|^{p-2} d_s u_N - |d_s \Phi_N|^{p-2} d_s \Phi_N \right) d_s \Phi_N \frac{dx dy}{|x-y|^n} = 0.$$

Proof. By the estimate (3.3) of Lemma 3.1, we obtain

$$\left| \int_{\mathbb{R}^{2n}} \sigma \left(|d_s u_N|^{p-2} d_s u_N - |d_s \Phi_N|^{p-2} d_s \Phi_N \right) d_s \Phi_N \frac{dx dy}{|x-y|^n} \right| \lesssim I,$$

where

$$I := \|\sigma\|_{L^\infty(\mathbb{R}^{2n})} \int_{\mathbb{R}^{2n}} (|d_s u_N| + |d_s \Phi_N|)^{p-2} |d_s u_N - d_s \Phi_N| |d_s \Phi_N| \frac{dx dy}{|x-y|^n}.$$

We divide the cases into $p \geq 2$ and $1 < p < 2$.

(i) For $p \geq 2$: The Hölder's inequality implies that

$$\begin{aligned} I &\lesssim \left(\| |d_s u_N| + |d_s \Phi_N| \|_{L^p(\mathbb{R}^{2n}, |x-y|^{-n})}^{p-2} \right) \\ &\quad \cdot \|d_s(u_N - \Phi_N)\|_{L^p(\mathbb{R}^{2n}, |x-y|^{-n})} \|d_s \Phi_N\|_{L^p(\mathbb{R}^{2n}, |x-y|^{-n})} \\ &\lesssim ([u_N]_{W^{s,p}(\mathbb{R}^n)} + [\Phi_N]_{W^{s,p}(\mathbb{R}^n)})^{p-2} [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)} [\Phi_N]_{W^{s,p}(\mathbb{R}^n)}. \end{aligned}$$

Since, $u_N \in W^{s,p}(\mathbb{R}^n)$ minimizes the fractional p -Dirichlet energy and $[\Phi_N]_{W^{s,p}(\mathbb{R}^n)} = 1$, Lemma 4.5 shows that this term goes to zero as $N \rightarrow \infty$.

(ii) For $1 < p < 2$: We obtain the same estimate from

$$\begin{aligned} &\int_{\mathbb{R}^{2n}} (|d_s u_N| + |d_s \Phi_N|)^{p-2} |d_s u_N - d_s \Phi_N| |d_s \Phi_N| \frac{dx dy}{|x-y|^n} \\ &\leq \int_{\mathbb{R}^{2n}} |d_s u_N - d_s \Phi_N|^{p-1} |d_s \Phi_N| \frac{dx dy}{|x-y|^n} \\ &\lesssim \underbrace{\|d_s(u_N - \Phi_N)\|_{L^p(\mathbb{R}^n, |x-y|^{-n})}^{p-1} \|d_s \Phi_N\|_{L^p(\mathbb{R}^n, |x-y|^{-n})}}_{\text{The Hölder's inequality.}} \\ &= [u_N - \Phi_N]_{W^{s,p}(\mathbb{R}^n)}^{p-1} [\Phi_N]_{W^{s,p}(\mathbb{R}^n)}. \end{aligned}$$

Here we used that $p-2 < 0$ implies that

$$(|d_s u_N| + |d_s \Phi_N|)^{p-2} \leq |d_s u_N - d_s \Phi_N|^{p-2}.$$

Arguing as in the case $p \geq 2$, we see that the last expression goes to zero as $N \rightarrow \infty$.

Hence, we can conclude the proof. \square

Lemma 4.7. *Let $\Omega \subset \mathbb{R}^n$ be a bounded open set, $0 < s < 1$, $1 < p < \infty$ and $x_0 \in W \Subset \Omega_e$. Assume that $(\Phi_N)_{N \in \mathbb{N}} \subset C_c^\infty(W)$ is a sequence satisfying the properties (i) – (iii) of Lemma 4.1 and let $\sigma: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ satisfy the uniform ellipticity condition (1.2). Then there holds*

$$\lim_{N \rightarrow \infty} \langle \Lambda_\sigma \Phi_N, \Phi_N \rangle = \lim_{N \rightarrow \infty} E_{s,p,\sigma}(\Phi_N).$$

Proof. As above, denote by $(u_N)_{N \in \mathbb{N}} \subset W^{s,p}(\mathbb{R}^n)$ the unique solutions to (4.6). Now by definition of the DN map Λ_σ and $u_N - \Phi_N \in \widetilde{W}^{s,p}(\Omega)$, there holds

$$\langle \Lambda_\sigma(\Phi_N), \Phi_N \rangle = E_{s,p,\sigma}(u_N).$$

Now the result directly follows from (4.5) and Proposition 4.6. \square

Finally, we can give the proof of Theorem 1.1.

Proof of Theorem 1.1. First note that such a sequence $(\Phi_N)_{N \in \mathbb{N}}$ exists by Lemma 4.1. By applying Lemma 4.3 and 4.7, we deduce

$$\sigma(x_0, x_0) = \lim_{N \rightarrow \infty} E_{s,p,\sigma}(\Phi_N) = \lim_{N \rightarrow \infty} \langle \Lambda_\sigma \Phi_N, \Phi_N \rangle.$$

Hence, we can conclude the proof. \square

5. PROOFS OF PROPOSITION 1.2, 1.3 AND 1.4

The proofs of Proposition 1.2, 1.3 and 1.4 can be regarded as corollaries of Theorem 1.1, and we give their proofs in the end of this work.

Proof of Proposition 1.2. Fix some $x_0 \in W$, we can choose a neighborhood V of x_0 such that $V \Subset \Omega_e$ and $V \subset W$. Then by Theorem 1.1 (with V in place of W), we deduce

$$\Sigma_j(x_0) = \sigma_j(x_0, x_0) = \lim_{N \rightarrow \infty} \langle \Lambda_{\sigma_j} \Phi_N, \Phi_N \rangle$$

for $j = 1, 2$. Since, the DN maps of σ_1 and σ_2 coincide for smooth functions compactly supported in W , we get $\Sigma_1(x_0) = \Sigma_2(x_0)$. As $x_0 \in W$ is arbitrary, we get $\Sigma_1 = \Sigma_2$ in W . \square

Proof of Proposition 1.3. Let $x_0 \in W$ and choose as above a neighborhood $V \Subset \Omega_e$ such that $x_0 \in V \subset W$. Then by Theorem 1.1 (with V in place of W), we have

$$\begin{aligned} & |\Sigma_1(x_0) - \Sigma_2(x_0)| \\ &= \lim_{N \rightarrow \infty} | \langle (\Lambda_{\sigma_1} - \Lambda_{\sigma_2}) \Phi_N, \Phi_N \rangle | \\ &\leq \limsup_{N \rightarrow \infty} \| \Lambda_{\sigma_1} - \Lambda_{\sigma_2} \|_{\widetilde{W}^{s,p}(V) \rightarrow (\widetilde{W}^{s,p}(V))^*} \| \Phi_N \|_{W^{s,p}(\mathbb{R}^n)} \| \Phi_N \|_{W^{s,p}(\mathbb{R}^n)} \\ &\leq \| \Lambda_{\sigma_1} - \Lambda_{\sigma_2} \|_{\widetilde{W}^{s,p}(W) \rightarrow (\widetilde{W}^{s,p}(W))^*}. \end{aligned}$$

In the last step, we used that by Theorem 1.1 the sequence $(\Phi_N)_{N \in \mathbb{N}} \subset C_c^\infty(V)$ satisfies $\| \Phi_N \|_{L^p(\mathbb{R}^n)} \rightarrow 0$ as $N \rightarrow \infty$ and $[\Phi_N]_{W^{s,p}(\mathbb{R}^n)} = 1$ for all $N \in \mathbb{N}$. Since, $x_0 \in W$ was arbitrary and the right hand side is independent of x_0 , we deduce

$$\| \Sigma_1 - \Sigma_2 \|_{L^\infty(W)} \leq \| \Lambda_{\sigma_1} - \Lambda_{\sigma_2} \|_{\widetilde{W}^{s,p}(W) \rightarrow (\widetilde{W}^{s,p}(W))^*}$$

and we can conclude the proof. \square

Proof of Proposition 1.4. Proposition 1.2 implies $\Sigma_1 = \Sigma_2$ on the nonempty open set $W \subset \Omega_e$. With the real-analyticity of Σ_j at hand, $j = 1, 2$, this immediately implies $\Sigma_1 = \Sigma_2$ in \mathbb{R}^n . \square

Acknowledgments. M.K. was supported by MATRICS grant (MTR/2019/001349) of SERB. Y.-H.L. was partially supported by NSTC (Taiwan) 113-2628-M-A49-003. P.Z. thanks Jesse Railo for fruitful discussions on the recovery of kernels on the diagonal in the linear case $p = 2$.

REFERENCES

- [AJ92] Gunnar Aronsson and Ulf Janfalk. On Hele-Shaw flow of power-law fluids. *European Journal of Applied Mathematics*, 3:343–366, December 1992.
- [AR06] Stanislav N. Antontsev and José F. Rodrigues. On stationary thermo-rheological viscous flows. *Annali dell'Universita di Ferrara*, 52(1):19–36, 2006.
- [Aro96] Gunnar Aronsson. On p -harmonic functions, convex duality and an asymptotic formula for injection mould filling. *European Journal of Applied Mathematics*, 7:417–437, October 1996.
- [BBM01] Jean Bourgain, Haim Brezis, and Petru Mironescu. Another look at Sobolev spaces. In *Optimal control and partial differential equations*, pages 439–455. IOS, Amsterdam, 2001.

- [BGU21] Sombuddha Bhattacharyya, Tuhin Ghosh, and Gunther Uhlmann. Inverse problems for the fractional-Laplacian with lower order non-local perturbations. *Trans. Amer. Math. Soc.*, 374(5):3053–3075, 2021.
- [BH22] Ali Behzadan and Michael Holst. Sobolev-slobodeckij spaces on compact manifolds, revisited. *Mathematics*, 10(3), 2022.
- [BHKS18] Tommi Brander, Bastian Harrach, Manas Kar, and Mikko Salo. Monotonicity and enclosure methods for the p -Laplace equation. *SIAM J. Appl. Math.*, 78(2):742–758, 2018.
- [BIK18] Tommi Brander, Joonas Ilmavirta, and Manas Kar. Superconductive and insulating inclusions for linear and non-linear conductivity equations. *Inverse Probl. Imaging*, 12(1):91–123, 2018.
- [BKS15] Tommi Brander, Manas Kar, and Mikko Salo. Enclosure method for the p -Laplace equation. *Inverse Problems*, 31(4):045001, 16, 2015.
- [BP12] Viorel Barbu and Teodor Precupanu. *Convexity and optimization in Banach spaces*. Springer Monographs in Mathematics. Springer, Dordrecht, fourth edition, 2012.
- [BPS16] Lorenzo Brasco, Enea Parini, and Marco Squassina. Stability of variational eigenvalues for the fractional p -Laplacian. *Discrete Contin. Dyn. Syst.*, 36(4):1813–1845, 2016.
- [Bra16] Tommi Brander. Calderón problem for the p -Laplacian: first order derivative of conductivity on the boundary. *Proc. Amer. Math. Soc.*, 144(1):177–189, 2016.
- [Bre11] Haim Brezis. *Functional analysis, Sobolev spaces and partial differential equations*. Universitext. Springer, New York, 2011.
- [Cal06] Alberto P Calderón. On an inverse boundary value problem. *Computational & Applied Mathematics*, 25(2-3):133–138, 2006.
- [CF24] Cătălin I Cârstea and Ali Feizmohammadi. Two uniqueness results in the inverse boundary value problem for the weighted p -Laplace equation. *arXiv preprint arXiv:2405.04123*, 2024.
- [CL19] Xinlin Cao and Hongyu Liu. Determining a fractional Helmholtz equation with unknown source and scattering potential. *Commun. Math. Sci.*, 17(7):1861–1876, 2019.
- [CLL19] Xinlin Cao, Yi-Hsuan Lin, and Hongyu Liu. Simultaneously recovering potentials and embedded obstacles for anisotropic fractional Schrödinger operators. *Inverse Probl. Imaging*, 13(1):197–210, 2019.
- [CLR20] Mihajlo Cekic, Yi-Hsuan Lin, and Angkana Rüland. The Calderón problem for the fractional Schrödinger equation with drift. *Cal. Var. Partial Differential Equations*, 59(91), 2020.
- [CMR21] Giovanni Covi, Keijo Mönkkönen, and Jesse Railo. Unique continuation property and Poincaré inequality for higher order fractional Laplacians with applications in inverse problems. *Inverse Probl. Imaging*, 15(4):641–681, 2021.
- [CMRU22] Giovanni Covi, Keijo Mönkkönen, Jesse Railo, and Gunther Uhlmann. The higher order fractional Calderón problem for linear local operators: Uniqueness. *Adv. Math.*, 399:Paper No. 108246, 2022.
- [CRTZ22] Giovanni Covi, Jesse Railo, Teemu Tyni, and Philipp Zimmermann. Stability estimates for the inverse fractional conductivity problem, 2022. arXiv: 2210.01875.
- [CRZ22] Giovanni Covi, Jesse Railo, and Philipp Zimmermann. The global inverse fractional conductivity problem. *arXiv:2204.04325*, 2022.
- [DNPV12] Eleonora Di Nezza, Giampiero Palatucci, and Enrico Valdinoci. Hitchhiker’s guide to the fractional Sobolev spaces. *Bull. Sci. Math.*, 136(5):521–573, 2012.
- [dTGCV21] Félix del Teso, David Gómez-Castro, and Juan Luis Vázquez. Three representations of the fractional p -laplacian: Semigroup, extension and balakrishnan formulas. *Fractional Calculus and Applied Analysis*, 24(4):966–1002, aug 2021.
- [FGKU21] Ali Feizmohammadi, Tuhin Ghosh, Katya Krupchyk, and Gunther Uhlmann. Fractional anisotropic Calderón problem on closed Riemannian manifolds. *arXiv:2112.03480*, 2021.
- [Gho22] Tuhin Ghosh. A non-local inverse problem with boundary response. *Rev. Mat. Iberoam.*, 38(6):2011–2032, 2022.
- [GK03] Adriana Garroni and Robert V Kohn. Some three-dimensional problems related to dielectric breakdown and polycrystal plasticity. *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences*, 459(2038):2613–2625, 2003.
- [GKS16] Chang-Yu Guo, Manas Kar, and Mikko Salo. Inverse problems for p -Laplace type equations under monotonicity assumptions. *Rend. Istit. Mat. Univ. Trieste*, 48:79–99, 2016.

- [GLX17] Tuhin Ghosh, Yi-Hsuan Lin, and Jingni Xiao. The Calderón problem for variable coefficients nonlocal elliptic operators. *Comm. Partial Differential Equations*, 42(12):1923–1961, 2017.
- [GM75] Roland Glowinski and Americo Marroco. Sur l’approximation, par éléments finis d’ordre un, et la résolution, par pénalisation-dualité, d’une classe de problèmes de Dirichlet non linéaires. *Rev. Française Automat. Informat. Recherche Opérationnelle Sér. Rouge Anal. Numér.*, 9(R-2):41–76, 1975.
- [GR03] Roland Glowinski and Jacques Rappaz. Approximation of a nonlinear elliptic problem arising in a non-Newtonian fluid flow model in glaciology. *ESAIM: Mathematical Modelling and Numerical Analysis*, 37:175–186, 1 2003.
- [GRSU20] Tuhin Ghosh, Angkana Rüland, Mikko Salo, and Gunther Uhlmann. Uniqueness and reconstruction for the fractional Calderón problem with a single measurement. *J. Funct. Anal.*, 279(1):108505, 42, 2020.
- [GSU20] Tuhin Ghosh, Mikko Salo, and Gunther Uhlmann. The Calderón problem for the fractional Schrödinger equation. *Anal. PDE*, 13(2):455–475, 2020.
- [GU21] Tuhin Ghosh and Gunther Uhlmann. The Calderón problem for nonlocal operators. *arXiv:2110.09265*, 2021.
- [HL19] Bastian Harrach and Yi-Hsuan Lin. Monotonicity-based inversion of the fractional Schrödinger equation I. Positive potentials. *SIAM J. Math. Anal.*, 51(4):3092–3111, 2019.
- [HL20] Bastian Harrach and Yi-Hsuan Lin. Monotonicity-based inversion of the fractional Schrödinger equation II. General potentials and stability. *SIAM J. Math. Anal.*, 52(1):402–436, 2020.
- [Idi08] Martín I. Idiart. The macroscopic behavior of power-law and ideally plastic materials with elliptical distribution of porosity. *Mechanics Research Communications*, 35(8):583–588, 2008.
- [KLL22] Minhyun Kim, Ki-Ahm Lee, and Se-Chan Lee. The Wiener criterion for nonlocal Dirichlet problems, 2022.
- [KLW22] Pu-Zhao Kow, Yi-Hsuan Lin, and Jenn-Nan Wang. The Calderón problem for the fractional wave equation: uniqueness and optimal stability. *SIAM J. Math. Anal.*, 54(3):3379–3419, 2022.
- [KRZ22] Manas Kar, Jesse Railo, and Philipp Zimmermann. The fractional p -biharmonic systems: optimal poincaré constants, unique continuation and inverse problems, 2022. *arXiv:2208.09528*.
- [KV84] Robert Kohn and Michael Vogelius. Determining conductivity by boundary measurements. *Comm. Pure Appl. Math.*, 37(3):289–298, 1984.
- [KW21] Manas Kar and Jenn-Nan Wang. Size estimates for the weighted p -Laplace equation with one measurement. *Discrete Contin. Dyn. Syst. Ser. B*, 26(4):2011–2024, 2021.
- [Lin22] Yi-Hsuan Lin. Monotonicity-based inversion of fractional semilinear elliptic equations with power type nonlinearities. *Calc. Var. Partial Differential Equations*, 61(5):Paper No. 188, 30, 2022.
- [Lin23] Yi-Hsuan Lin. Determining both leading coefficient and source in a nonlocal elliptic equation. *arXiv preprint arXiv:2312.15607*, 2023.
- [LK98] Ohad Levy and Robert V. Kohn. Duality relations for non-Ohmic composites, with applications to behavior near percolation. *Journal of Statistical Physics*, 90(1–2):159–189, 1998.
- [LL22a] Ru-Yu Lai and Yi-Hsuan Lin. Inverse problems for fractional semilinear elliptic equations. *Nonlinear Anal.*, 216:Paper No. 112699, 21, 2022.
- [LL22b] Yi-Hsuan Lin and Hongyu Liu. Inverse problems for fractional equations with a minimal number of measurements. *arXiv:2203.03010*, 2022.
- [LLR20] Ru-Yu Lai, Yi-Hsuan Lin, and Angkana Rüland. The Calderón problem for a space-time fractional parabolic equation. *SIAM J. Math. Anal.*, 52(3):2655–2688, 2020.
- [LLU22] Ching-Lung Lin, Yi-Hsuan Lin, and Gunther Uhlmann. The Calderón problem for nonlocal parabolic operators. *arXiv:2209.11157*, 2022.
- [LRZ22] Yi-Hsuan Lin, Jesse Railo, and Philipp Zimmermann. The Calderón problem for a nonlocal diffusion equation with time-dependent coefficients, 2022. *arXiv:2211.07781*.
- [MS18] Katarzyna Mazowiecka and Armin Schikorra. Fractional div-curl quantities and applications to nonlocal geometric equations. *Journal of Functional Analysis*, 275(1):1–44, jul 2018.
- [RO16] Xavier Ros-Oton. Nonlocal elliptic equations in bounded domains: a survey. *Publ. Mat.*, 60(1):3–26, 2016.
- [RS18] Angkana Rüland and Mikko Salo. Exponential instability in the fractional Calderón problem. *Inverse Problems*, 34(4):045003, 2018.

- [RS20] Angkana Rüland and Mikko Salo. The fractional Calderón problem: low regularity and stability. *Nonlinear Anal.*, 193:111529, 56, 2020.
- [RZ22a] Jesse Railo and Philipp Zimmermann. Counterexamples to uniqueness in the inverse fractional conductivity problem with partial data. *Inverse Probl. Imaging (to appear)*, 2022. arXiv:2203.02442.
- [RZ22b] Jesse Railo and Philipp Zimmermann. Fractional Calderón problems and Poincaré inequalities on unbounded domains. *J. Spectr. Theory (to appear)*, 2022. arXiv:2203.02425.
- [RZ22c] Jesse Railo and Philipp Zimmermann. Low regularity theory for the inverse fractional conductivity problem. *arXiv:2208.11465*, 2022.
- [Sim78] Jacques Simon. Régularité de la solution d'une équation non linéaire dans \mathbf{R}^N . In *Journées d'Analyse Non Linéaire (Proc. Conf., Besançon, 1977)*, volume 665 of *Lecture Notes in Math.*, pages 205–227. Springer, Berlin, 1978.
- [Str08] Michael Struwe. *Variational methods*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, fourth edition, 2008. Applications to nonlinear partial differential equations and Hamiltonian systems.
- [SU87] John Sylvester and Gunther Uhlmann. A global uniqueness theorem for an inverse boundary value problem. *Ann. of Math. (2)*, 125(1):153–169, 1987.
- [SZ12] Mikko Salo and Xiao Zhong. An inverse problem for the p -Laplacian: Boundary determination. *SIAM J. Math. Anal.*, 44(4):2474–2495, March 2012.
- [TW94a] DRS Talbot and John Raymond Willis. Upper and lower bounds for the overall properties of a nonlinear composite dielectric. I. Random microgeometry. *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences*, 447(1930):365–384, 1994. With second part [TW94b].
- [TW94b] DRS Talbot and John Raymond Willis. Upper and lower bounds for the overall properties of a nonlinear composite dielectric. II. Periodic microgeometry. *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences*, 447(1930):385–396, 1994. With first part [TW94a].
- [Wol07] Thomas H. Wolff. Gap series constructions for the p -Laplacian. *Journal d'Analyse Mathématique*, 102(1):371–394, August 2007. Preprint written in 1984.

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) BHOPAL, INDIA
Email address: manas@iiserb.ac.in

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL YANG MING CHIAO TUNG UNIVERSITY,
 HSINCHU, TAIWAN
Email address: yihuanlin3@gmail.com

DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, ZÜRICH, SWITZERLAND
Email address: philipp.zimmermann@math.ethz.ch