

# THE FRACTIONAL ANISOTROPIC CALDERÓN PROBLEM FOR A NONLOCAL PARABOLIC EQUATION ON CLOSED RIEMANNIAN MANIFOLDS

YI-HSUAN LIN

ABSTRACT. We consider the fractional anisotropic Calderón problem for the nonlocal parabolic equation  $(\partial_t - \Delta_g)^s u = f$  ( $0 < s < 1$ ) on closed Riemannian manifolds. More concretely, we can determine the Riemannian manifold  $(M, g)$  up to isometry by using the local source-to-solution map in an arbitrarily small open cylinder in the spacetime domain. This can be regarded as a nonlocal analog of the anisotropic Calderón problem in the parabolic setting. We also study several useful properties for nonlocal parabolic operators by using comprehensive spectrum analysis with semigroup theory.

## CONTENTS

|   |    |
|---|----|
| 1. Introduction   | 1  |
| 2. The nonlocal parabolic operator on closed Riemannian manifolds | 4  |
| 3. The well-posedness   | 7  |
| 4. The Calderón problem   | 8  |
| 4.1. Auxiliary lemmas   | 8  |
| 4.2. Proof of Theorem 1.1   | 10 |
| References  | 13 |

## 1. INTRODUCTION

In 1980, A. P. Calderón [Cal06] proposed a problem: Can the (anisotropic) conductivity be recovered in a medium by using its voltage and current measurements on the boundary? This problem is often called the *anisotropic Calderón problem*, which can be reformulated in the geometrical setting, see [LU89]. In other words, can one recover  $(M, g)$  by using the boundary measurements (usually, researchers utilize either the Dirichlet-to-Neumann map or the Neumann-to-Dirichlet map)? In the works [LU89, LU01, LLS20], the authors studied the anisotropic Calderón problem on real-analytic manifolds with real-analytic metrics. However, without using analyticity assumptions, the anisotropic Calderón problem is a well-known long-standing open problem in the inverse problems community for the spatial dimension  $n \geq 3$ .

In the works [FGKU24, Fei24, FKU24], the authors investigated the fractional anisotropic Calderón problem on closed connected smooth Riemannian manifolds, which they used the local source-to-solution map as measurement so that they can determine the manifold up to isometry. This type of result is different from that of their local counterparts. On the one hand, the fractional anisotropic Calderón

---

2020 *Mathematics Subject Classification.* 26A33, 35R30.

*Key words and phrases.* Anisotropic Calderón's problem, nonlocal parabolic operators, closed Riemannian manifolds, heat kernel.

problem can be resolved. On the other hand, the result holds for any spatial dimension  $n \geq 2$ , without additional analytic assumptions on manifolds  $(M, g)$ . The study of fractional inverse problems has been an attractive research field. Thanks to the pioneering work [GSU20], the authors proved two important materials: The strong uniqueness and Runge approximation. Motivated by the work [GSU20], there are many articles studied fractional type inverse problems, such as [GRSU20, CLR20, CLL19, CMR21, CMRU22, GLX17, CGRU23, GU21, HL19, HL20, LL22, LL23, LLR20, LLU22, LLU23, Lin22, Lin23, Rül23, RS18, RS20, KLV22, KRZ23, KLZ24, LZ24, LZ23, LTZ24a, LTZ24b, LNZZ24, FL24] and references therein. It is worth to mentioning that an entanglement principle for the fractional Laplacian has been found in [FKU24, FL24] for both compact and non-compact manifolds.

The mathematical model in this paper is given as follows. Let  $(M, g)$  be a smooth closed Riemannian manifold of dimension  $\dim M \geq 2$ , and  $-\Delta_g$  be the positive Laplace-Beltrami operator defined on  $M$ , i.e.,  $g = g(x) \in C^\infty(M)$  satisfies the classical ellipticity condition (we do not address more in this paper). A closed manifold is compact without boundary. Given  $s \in (0, 1)$  and  $T > 0$ , let

$$\mathcal{H} := \partial_t - \Delta_g,$$

and we consider the fractional parabolic equation

$$(1.1) \quad \begin{cases} \mathcal{H}^s u = f & \text{in } M \times (-T, T) \\ u = 0 & \text{in } M \times \{t \leq -T\}, \end{cases}$$

where the fractional parabolic operator  $\mathcal{H}^s$  can be defined by the spectrum theory (see Section 2). The well-posedness of (1.1) holds for any  $f$  in a suitable function space with the compatibility condition

$$(1.2) \quad f = 0 \text{ in } M \times \{t \leq -T\}.$$

Let  $\mathcal{O} \subset M$  be an open set, by the well-posedness of (1.1), we can define the local source-to-solution map  $S_{M,g,\mathcal{O}}$  of (1.1) via

$$(1.3) \quad S_{M,g,\mathcal{O}} : C_c^\infty(\mathcal{O} \times (-T, T)) \ni f \mapsto u^f|_{\mathcal{O} \times (-T, T)} \in \mathbf{H}^s(\mathcal{O} \times (-T, T)),$$

where  $u^f \in \mathbf{H}^s(M \times \mathbb{R})$  is the solution to (1.1). Here  $\mathbf{H}^s(M \times \mathbb{R})$  is a suitable function space given Section 2.

Recall that  $-\Delta_g$  is a nonnegative linear operator on the Hilbert space  $L^2(M)$ . Since  $M$  is a connected closed smooth Riemannian, it is known that  $-\Delta_g$  possesses a discrete spectrum, which can be denoted as an eigenpairs  $\{(\lambda_k, \phi_k)\}_{k \geq 0}$ , such that  $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots \nearrow \infty$  (up to normalization) and  $\{\phi_k\}_{k \geq 0}$  is an orthonormal basis in  $L^2(M)$ . Notice that these eigenvalues  $\lambda_k$  may not be distinct. The zeroth eigenfunction  $\phi_0 \equiv 1$  in  $M$ . This implies that

$$(1.4) \quad (\phi_k, 1)_{L^2(M)} = (\phi_k, \phi_0)_{L^2(M)} = 0, \quad \text{for all } k \in \mathbb{N}.$$

Furthermore, it is known that the fractional Laplace-Beltrami operator can be defined by

$$(-\Delta_g)^s u(x) = \sum_{k=1}^{\infty} \lambda_k^s (u, \phi_k)_{L^2(M)} \phi_k(x) \text{ for } x \in M,$$

where we use  $\lambda_0 = 0$ , and  $(\varphi, \psi)_{L^2(D)} := \int_D \varphi \bar{\psi} \, dV_g$ , for all  $\varphi, \psi \in L^2(M)$ . Here  $D$  can be any subset of  $M$ , and  $dV_g$  is the Riemannian volume element of  $(M, g)$  with  $\bar{\psi}$  being the complex conjugate of the (complex) function  $\psi$ . Similar  $L^2$  integral formula notation will be utilized for any subsets in the space-time domain  $M \times \mathbb{R}$ . Note that  $((-\Delta_g)^s u, 1)_{L^2(M)} = 0$ , for any  $u \in \text{Dom}((-\Delta_g)^s)$ , where we used (1.4) and  $\text{Dom}((-\Delta_g)^s) := \{u \in L^2(M) : \sum_{k=0}^{\infty} \lambda_k^{2s} |(u, \phi_k)_{L^2(M)}|^2 < \infty\}$ .

Motivated by the above observations, the nonlocal parabolic operator  $\mathcal{H}^s$  can be defined spectrally. More precisely, given any function  $u = u(x, t) \in L^2(M \times \mathbb{R})$ , one can write

$$u(x, t) = \sum_{k=0}^{\infty} u_k(t) \phi_k(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \sum_{k=0}^{\infty} \widehat{u}_k(\rho) \phi_k(x) e^{it\rho} dt,$$

where  $i = \sqrt{-1}$ . Here  $u_k(t)$  and  $\widehat{u}_k(\rho)$  (the time-Fourier transform of  $u_k(t)$ ) are given by

$$(1.5) \quad u_k(t) := (u(\cdot, t), \phi_k(\cdot))_{L^2(M)} \quad \text{and} \quad \widehat{u}_k(\rho) := \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-i\rho t} u_k(t) dt,$$

respectively, for  $k \geq 0$ . The fractional parabolic operator  $\mathcal{H}^s$  is (formally) defined by the pairing

$$\langle \mathcal{H}^s u, v \rangle = \int_{\mathbb{R}} \sum_{k=0}^{\infty} (i\rho + \lambda_k)^s \widehat{u}_k(\rho) \overline{\widehat{v}_k(\rho)} d\rho,$$

where we will justify the above pairing in Section 2 under appropriate sense with suitable function spaces. We will introduce more properties of the operator  $\mathcal{H}^s$  in Section 2. Our main result to determine  $(M, g)$  (up to isometry) by using the local source-to-solution map (1.3) is stated in the forthcoming theorem:

**Theorem 1.1.** *Given  $s \in (0, 1)$ , let  $(M_1, g_1)$  and  $(M_2, g_2)$  be closed connected smooth Riemannian manifolds of  $\dim M_1 = \dim M_2 \geq 2$ . Let  $\mathcal{O}_j \subset M_j$  be a nonempty open sets (for  $j = 1, 2$ ) such that*

$$(1.6) \quad (\mathcal{O}_1, g_1|_{\mathcal{O}_1}) = (\mathcal{O}_2, g_2|_{\mathcal{O}_2}) := (\mathcal{O}, g).$$

Suppose that

$$(1.7) \quad S_{M_1, g_1, \mathcal{O}_1}(f) = S_{M_2, g_2, \mathcal{O}_2}(f), \quad \text{for any } f \in C_c^\infty(\mathcal{O} \times (-T, T)),$$

where  $S_{M_j, g_j, \mathcal{O}_j}$  denotes the local source-to-solution map of

$$\begin{cases} (\partial_t - \Delta_{g_j})^s u_j = f & \text{in } M \times (-T, T), \\ u_j = 0 & \text{in } M \times \{t \leq -T\}, \end{cases}$$

for  $j = 1, 2$ . Then there exists a diffeomorphism  $\Phi : M_1 \rightarrow M_2$  such that  $\Phi^* g_2 = g_1$  in  $M_1$   $\Phi = \text{Id}$  (the identity map) on  $\mathcal{O}$ .

The preceding theorem was demonstrated in [FGKU24] for the elliptic case. Since  $(\partial_t - \Delta_g)^s$  is a space-like nonlocal operator (as  $g = g(x)$ ), we could expect a similar result for this nonlocal parabolic operator.

**Strategy.** To prove Theorem 1.1, our strategy is to demonstrate that the conditions (1.6) and (1.7) can be used to deduce the identity

$$(1.8) \quad e^{-\tau \mathcal{L}_1}(x, z) = e^{-\tau \mathcal{L}_2}(x, z), \quad \text{for } x, z \in \mathcal{O} \text{ and } \tau > 0,$$

where  $\mathcal{L}_j := -\Delta_{g_j}$ , for  $j = 1, 2$ . If the condition (1.8) holds, then one can apply [FGKU24, Theorem 1.5] to conclude Theorem 1.1 as we wish. For this purpose, we need to analyze the function  $e^{-\tau \mathcal{H}} f$  for any  $f \in C_c^\infty(\mathcal{O} \times (-T, T))$  carefully (see Sections 2 and 4).

**Organization of the article.** The paper is organized as follows. In Section 2, we provide a rigorous definition of the nonlocal parabolic operator via the spectrum theory and related function spaces will be introduced. Furthermore, we demonstrate that the future data  $u|_{\{t \geq T\}}$  of the solution  $u$  to (1.1) will not affect the solution  $u|_{\{-T \leq t \leq T\}}$ . In Section 3, we prove the well-posedness of (1.1), which also shows

that the operator  $\mathcal{H}^s$  is a space-like nonlocal operator. Finally, the proof of Theorem 1.1 will be given in Section 4, which can be derived by showing the conditions (1.6) and (1.7) ensure the identity (1.8).

## 2. THE NONLOCAL PARABOLIC OPERATOR ON CLOSED RIEMANNIAN MANIFOLDS

Given  $a \in \mathbb{R}$ , let us denote

$$\mathbf{H}^a(M \times \mathbb{R}) := \{u \in L^2(M \times \mathbb{R}) : \|u\|_{\mathbf{H}^a(M \times \mathbb{R})} < \infty\},$$

where

$$\|u\|_{\mathbf{H}^a(M \times \mathbb{R})} = \left( \int_{\mathbb{R}} \sum_{k=0}^{\infty} \left(1 + |\rho + \lambda_k|^2\right)^{a/2} |\widehat{u}_k(\rho)|^2 d\rho \right)^{1/2}.$$

Here

$$|\rho + \lambda_k|^2 = \rho^2 + \lambda_k^2, \text{ for } k \in \mathbb{N} \cup \{0\}.$$

It is obvious that  $\mathbf{H}^s(M \times \mathbb{R}) \subset L^2(M \times \mathbb{R})$ . Moreover, for any open set  $O \subset M \times \mathbb{R}$ , let us denote

$$\begin{aligned} \mathbf{H}^a(O) &:= \{u|_O : u \in \mathbf{H}^s(M \times \mathbb{R})\}, \\ \widetilde{\mathbf{H}}^a(O) &:= \text{closure of } C_c^\infty(O) \text{ in } \mathbf{H}^a(M \times \mathbb{R}). \end{aligned}$$

By the duality, there holds that

$$(\mathbf{H}^a(O))^* = \widetilde{\mathbf{H}}^{-a}(O) \quad \text{and} \quad (\widetilde{\mathbf{H}}^a(O))^* = \mathbf{H}^{-a}(O).$$

We also write

$$\mathbf{H}_F^s = \{u \in \mathbf{H}^s(M \times \mathbb{R}) : \text{supp}(u) \subset F\},$$

for any closed set  $F \subset M \times \mathbb{R}$ .

Let us first review several useful properties for the nonlocal operator  $\mathcal{H}^s$ . Given  $s \in (0, 1)$ , let us define the domain of  $\mathcal{H}^s$  by

$$\text{Dom}(\mathcal{H}^s) = \mathbf{H}^{2s}(M \times \mathbb{R})$$

where Here  $(\lambda_k)_{k \geq 0}$  is the eigenvalue of  $-\Delta_g$  in  $M$  introduced in the previous section. We denote  $\mathbf{H}^{-s}(M \times \mathbb{R})$  as the dual space of  $\mathbf{H}^s(M \times \mathbb{R})$ , and it is not hard to check that

$$(2.1) \quad \begin{aligned} \mathcal{H}^s &: \mathbf{H}^s(M \times \mathbb{R}) \rightarrow \mathbf{H}^{-s}(M \times \mathbb{R}), \\ \mathcal{H}^s &: \mathbf{H}^{s/2}(M \times \mathbb{R}) \rightarrow L^2(M \times \mathbb{R}). \end{aligned}$$

The first relation can be seen by using the classical pairing argument. More concretely, one has that

$$(2.2) \quad (\mathcal{H}^s u, v)_{L^2(M \times \mathbb{R})} = \int_{\mathbb{R}} \sum_{k=0}^{\infty} (i\rho + \lambda_k)^s \widehat{u}_k(\rho) \overline{\widehat{v}_k(\rho)} d\rho,$$

for any  $u, v \in C^\infty(M \times \mathbb{R})$ . Then a density argument ( $C^\infty(M \times \mathbb{R})$  is dense in  $\mathbf{H}^s(M \times \mathbb{R})$  with respect to the norm  $\|\cdot\|_{\mathbf{H}^s}$ ) yields that (2.2) holds, so that the mapping property (2.1) is valid in the following form

$$\langle \mathcal{H}^s u, v \rangle_{\mathbf{H}^{-s} \times \mathbf{H}^s} = \int_{\mathbb{R}} \sum_{k=0}^{\infty} (i\rho + \lambda_k)^s \widehat{u}_k(\rho) \overline{\widehat{v}_k(\rho)} d\rho$$

and we use  $\|u\|_{\mathbf{H}^s} = \|u\|_{\mathbf{H}^s(M \times \mathbb{R})}$  for the simplicity. This shows the first property in (2.1). For the second property, it is easy to see by using the definition of  $\mathcal{H}^s$ . More generally, one can also derive that  $\mathcal{H}^b$  satisfies

$$\mathcal{H}^b : \mathbf{H}^a(M \times \mathbb{R}) \rightarrow \mathbf{H}^{a-2b}(M \times \mathbb{R}),$$

for any  $a \in \mathbb{R}$  and  $b \geq 0$ .

On the other hand, we have the following integration-by-parts formula that

$$(2.3) \quad (\mathcal{H}^s u, v)_{\mathbf{H}^{-s} \times \mathbf{H}^s} = (\mathcal{H}^{s/2} u, \mathcal{H}_*^{s/2} v)_{L^2(M \times \mathbb{R})},$$

where  $\mathcal{H}_*^s := (-\partial_t - \Delta_g)^s$  denotes the adjoint operator of  $\mathcal{H}^s$ , which is also a time-reversal operator of  $\mathcal{H}^s$ . Similar to the definition of  $\mathcal{H}^s$ , one can define  $\mathcal{H}_*^s$  via

$$\langle \mathcal{H}_*^s v, u \rangle_{\mathbf{H}^{-s} \times \mathbf{H}^s} = \int_{\mathbb{R}} \sum_{k=0}^{\infty} (-i\rho + \lambda_k)^s \widehat{v}_k(\rho) \overline{\widehat{u}_k(\rho)} d\rho,$$

where  $\widehat{u}_k(\rho)$  and  $\widehat{v}_k(\rho)$  are given by (1.5).

Recall that  $(\phi_k)_{k \geq 0}$  is an orthonormal basis in  $L^2(M)$ , and we can write the semigroup  $\{e^{-\tau \mathcal{L}}\}_{\tau \geq 0}$  generated by  $\mathcal{L} := -\Delta_g$  on  $M$  as

$$(2.4) \quad e^{-\tau \mathcal{L}} \varphi(x) = \sum_{k=0}^{\infty} e^{-\tau \lambda_k} (\varphi, \phi_k)_{L^2(M)} \phi_k = \int_M e^{-\tau \mathcal{L}}(x, z) \varphi(z) dz,$$

for any  $\varphi \in L^2(M)$  and for  $k \geq 0$ . Here  $e^{-\tau \mathcal{L}}(x, z)$  is the heat kernel of  $\partial_t - \Delta_g = \partial_t + \mathcal{L}$  on  $M$ , which is symmetric and nonnegative that  $e^{-\tau \mathcal{L}}(x, z) = e^{-\tau \mathcal{L}}(z, x) \geq 0$ , for all  $x, z \in M$ . Moreover,  $e^{-\tau \mathcal{L}}(x, z)$  possesses the Gaussian upper bound

$$(2.5) \quad e^{-\tau \mathcal{L}}(x, z) \leq \frac{C}{\tau^{n/2}} e^{-\frac{|x-z|^2}{c\tau}}, \text{ for all } x, z \in M \text{ and } \tau > 0,$$

where  $c, C > 0$  are some positive constants (cf. [SY94, Theorem 4.7, page 17]). On the other hand, via the condition (1.4), observe that

$$(2.6) \quad e^{-\tau \mathcal{L}} 1(x) = 1, \text{ for all } x \in M \text{ and } \tau > 0,$$

whenever  $(M, g)$  is a closed Riemannian manifold.

Since  $\partial_t$  and  $-\Delta_g$  are commutable, we have  $e^{-\tau \mathcal{H}} = e^{-\tau \mathcal{L}} \circ e^{-\tau \partial_t}$  and

(2.7)

$$e^{-\tau \mathcal{H}} u(x, t) = e^{-\tau \mathcal{L}} (e^{-\tau \partial_t} u) = e^{-\tau \mathcal{L}} (u(\cdot, t - \tau))(x) = \int_M e^{-\tau \mathcal{L}}(x, z) u(z, t - \tau) dz$$

in the sense that

$$\begin{aligned} (e^{-\tau \mathcal{H}} u, v)_{L^2(M \times \mathbb{R})} &= \int_{\mathbb{R}} \sum_{k=0}^{\infty} e^{-\tau(i\rho + \lambda_k)} \widehat{u}_k(\rho) \overline{\widehat{v}_k(\rho)} d\rho \\ &= \int_{\mathbb{R}} \sum_{k=0}^{\infty} e^{-\tau \lambda_k} u_k(t - \tau) v_k(t) dt \\ &= \int_{\mathbb{R}} \int_M \int_M e^{-\tau \mathcal{L}}(x, z) u(z, t - \tau) v(x, t) dz dx dt. \end{aligned}$$

Recalling that  $(M, g)$  is a closed connected smooth Riemannian manifold, by using the formula (2.2), it is not hard to see

$$(2.8) \quad \|e^{-\tau \mathcal{H}} u\|_{L^2(M \times \mathbb{R})} \leq e^{-c_0 \tau} \|u\|_{L^2(M \times \mathbb{R})}, \quad \text{for } \tau \geq 0,$$

for some constant  $c_0 > 0$ .

With the above expressions and formulae, we can show an alternative formulation used to characterize  $\mathcal{H}^s$  by the heat semigroup method.

**Lemma 2.1.** *Given  $s \in (0, 1)$ , for any  $u \in \mathbf{H}^s(M \times \mathbb{R})$ , there holds that*

$$(2.9) \quad \mathcal{H}^s u = \frac{1}{\Gamma(-s)} \int_0^{\infty} (e^{-\tau \mathcal{H}} u - u) \frac{d\tau}{\tau^{1+s}}$$

in the sense that

$$\langle \mathcal{H}^s u, v \rangle_{\mathbf{H}^{-s} \times \mathbf{H}^s} = \frac{1}{\Gamma(-s)} \int_0^{\infty} ((e^{-\tau \mathcal{H}} u, v)_{L^2(M \times \mathbb{R})} - (u, v)_{L^2(M \times \mathbb{R})}) \frac{d\tau}{\tau^{1+s}},$$

for any  $v \in \mathbf{H}^s(M \times \mathbb{R})$ .

*Proof.* By using the formula

$$(i\rho + \lambda_k)^s = \frac{1}{\Gamma(-s)} \int_0^\infty (e^{-\tau(i\rho + \lambda_k)} - 1) \frac{d\tau}{\tau^{1+s}},$$

where the integral converges absolutely (see the proof of [BDLCS21, Lemma 2.1]). Combining with the formula (2.2), there holds that

$$\langle \mathcal{H}^s u, v \rangle_{\mathbf{H}^{-s} \times \mathbf{H}^s} = \int_{\mathbb{R}} \sum_{k=0}^{\infty} \left[ \frac{1}{\Gamma(-s)} \int_0^\infty (e^{-\tau(i\rho + \lambda_k)} - 1) \frac{d\tau}{\tau^{1+s}} \right] \widehat{u}_k(\rho) \overline{\widehat{v}_k(\rho)} d\rho,$$

which proves the assertion.  $\square$

Let us refer readers to the *Balakrishnan formula* (see [Sam02, equation (9.63), page 285]), which can be used to define the general fractional order of operators. Last but not least, we can derive an integral representation formula of  $\mathcal{H}^s$ .

**Proposition 2.2** (Representation formula). *We have the integral formula of  $\mathcal{H}^s$  by*

$$(2.10) \quad \mathcal{H}^s u(x, t) = \int_0^\infty \int_M \mathcal{K}_s(x, z; \tau) (u(z, t - \tau) - u(x, t)) dz d\tau$$

which belong to  $\mathbf{H}^{-s}(M \times \mathbb{R})$ , for any  $u \in \mathbf{H}^s(M \times \mathbb{R})$ , where

$$(2.11) \quad \mathcal{K}_s(x, z; \tau) := \frac{1}{\Gamma(-s)} \frac{e^{-\tau \mathcal{L}}(x, z)}{\tau^{1+s}}, \text{ for } x, z \in M \text{ and } \tau > 0.$$

*Proof.* Note that from the formulae (2.4) and (2.6), then there holds that

$$(2.12) \quad 1 = e^{-\tau \mathcal{L}} 1(x) = \int_M e^{-\tau \mathcal{L}}(x, z) dz, \text{ for all } x \in M.$$

Now, by using (2.7) and (2.9), we have

$$\begin{aligned} & \mathcal{H}^s u(x, t) \\ &= \frac{1}{\Gamma(-s)} \int_0^\infty \left( \int_M e^{-\tau \mathcal{L}}(x, z) u(z, t - \tau) dz - u(x, t) \right) \frac{d\tau}{\tau^{1+s}} \\ &= \frac{1}{\Gamma(-s)} \int_0^\infty \left( \int_M e^{-\tau \mathcal{L}}(x, z) u(z, t - \tau) dz - \underbrace{\left( \int_M e^{-\tau \mathcal{L}}(x, z) dz \right)}_{=1 \text{ by (2.12)}} u(x, t) \right) \frac{d\tau}{\tau^{1+s}} \\ &= \frac{1}{\Gamma(-s)} \int_0^\infty \int_M e^{-\tau \mathcal{L}}(x, z) (u(z, t - \tau) - u(x, t)) dz \frac{d\tau}{\tau^{1+s}}, \end{aligned}$$

where we use the Fubini theorem in the last identity. Finally, using (2.11), this proves the assertion.  $\square$

**Remark 2.3.** *If  $u$  is time-independent, i.e.,  $u(x, t) \equiv u(x)$ , it is easy to see that  $\mathcal{H}^s u(x) = (-\Delta_g)^s u(x)$  by using the formula (2.10). However, from the initial data of the problem (1.1), one can not simply take the solution  $u(x, t)$  as a time-independent function so that one can apply the existing results given in [FGKU24] to show Theorem 1.1.*

Motivated by [LLR20], we have the following lemma.

**Lemma 2.4.** *Let  $\chi_A(t)$  be the characteristic function of  $A \subset \mathbb{R}$ , for any  $u \in \mathbf{H}^s(M \times \mathbb{R})$ , then there holds that*

- (i)  $\chi_A(t)u(x, t) \in \mathbf{H}^s(M \times \mathbb{R})$ , for any interval  $A \subset \mathbb{R}$ .
- (ii)  $\mathcal{H}^s u(x, t) = \mathcal{H}^s (\chi_{(-\infty, T]}(t)u(x, t))$  for  $(x, t) \in \Omega \times (-T, T)$ .

*Proof.* Since  $\chi_A(t)$  is a space-independent function, the first statement follows by utilizing a similar approach given in [LLR20, Section 3]. For the second statement, note that  $\chi_{(-\infty, T]}(t)u \in \mathbf{H}^s(M \times \mathbb{R})$  provided that  $u \in \mathbf{H}^s(M \times \mathbb{R})$ . Then the identity holds by using the integral representation (2.10). More specifically, one may compute that

$$\begin{aligned} & \mathcal{H}^s u(x, t) \\ &= \int_0^\infty \int_M \mathcal{K}_s(x, z; \tau) (u(z, t - \tau) - u(x, t)) dz d\tau \\ &= \int_0^\infty \int_M \mathcal{K}_s(x, z; \tau) ((\chi_{(-\infty, T]}(t)u)(z, t - \tau) - (\chi_{(-\infty, T]}(t)u)(x, t)) dz d\tau \end{aligned}$$

for  $(x, t) \in M \times (-T, T)$ , and  $u \in \mathbf{H}^s(M \times \mathbb{R})$ , and the last identity holds since  $\tau \in (0, \infty)$ . This completes the proof.  $\square$

The above lemma states that the value  $\mathcal{H}^s u|_{M \times (-T, T)}$  will not be affected by the information  $u|_{\{t \geq T\}}$ . In other words, the value  $\mathcal{H}^s u(x, t)$  depends only on the past, not the future. Similar properties as in Lemma 2.4 hold for the adjoint operator  $\mathcal{H}_*^s$ , and the value of  $\mathcal{H}_*^s u|_{M \times [-T, T]}$  will not be affected by the past time  $\{t \leq -T\}$ .

### 3. THE WELL-POSEDNESS

We next turn to study the well-posedness of (1.1). As explained in [LLR20], the operator  $\mathcal{H}^s$  is a space-time coupled operator, which is the space-like nonlocal operator. Thus, combining with the above relation (2.3), one can define the bilinear form associated with (1.1) via

$$B_g(u, v) := (\mathcal{H}^{s/2} u, \mathcal{H}_*^{s/2} v)_{L^2(M \times \mathbb{R})}.$$

then similar to the arguments as in [LLR20, BS24], one can check that  $B_g(\cdot, \cdot)$  is bounded and coercive in the space  $\mathbf{H}^s(M \times \mathbb{R})$ . It is worth mentioning that this argument was first found in the work [LLR20], which also demonstrates that  $\mathcal{H}^s$  is a space-like nonlocal operator.

**Lemma 3.1** (Well-posedness). *Let  $(M, g)$  be a closed connected smooth Riemannian manifold with  $\dim M \geq 2$ . Then given  $f \in \mathbf{H}^{-s}(M \times \mathbb{R})$  fulfilling (1.2), there exists a unique solution  $u \in H^s(M \times \mathbb{R})$  solving (1.1).*

*Proof.* Notice that even  $\mathcal{H}^s$  is a fractional parabolic operator, we will use the Lax-Milgram theorem to prove the result. To simplify the notation, let us denote

$$u_T(x, t) = \chi_{[-T, T]}(t)u(x, t),$$

which belongs to  $\mathbf{H}^s(M \times \mathbb{R})$  as  $u \in \mathbf{H}^s(M \times \mathbb{R})$  (see Lemma 2.4). On one hand, for the boundedness, by using the Hölder inequality, we have

$$\begin{aligned} & |B_g(u_T, v)| \\ &= \left| \int_{\mathbb{R}} \sum_{k=0}^{\infty} (i\rho + \lambda_k)^s \widehat{u_{T,k}}(\rho) \overline{\widehat{v_k}(\rho)} d\rho \right| \\ (3.1) \quad & \leq \left( \int_{\mathbb{R}} \sum_{k=0}^{\infty} |i\rho + \lambda_k|^s |\widehat{u_{T,k}}(\rho)|^2 d\rho \right)^{1/2} \left( \int_{\mathbb{R}} \sum_{k=0}^{\infty} |i\rho + \lambda_k|^s |\widehat{v_k}(\rho)|^2 d\rho \right)^{1/2} \\ & \leq \|u_T\|_{\mathbf{H}^s(M \times \mathbb{R})} \|v\|_{\mathbf{H}^s(M \times \mathbb{R})}, \end{aligned}$$

for any  $v \in \mathbf{H}_{M \times [-T, T]}^s$ , where  $u_{T,k}$  and  $\widehat{u_{T,k}}$  are the functions given by (1.5) when the function  $u$  is replaced by  $u_T$ .

On the other hand, for the coercive, we can see that

$$\begin{aligned}
(3.2) \quad B_g(u_T, u_T) &= \int_{\mathbb{R}} \sum_{k=0}^{\infty} (i\rho + \lambda_k)^s |\widehat{u_{T,k}}(\rho)|^2 d\rho \\
&= \int_{\mathbb{R}} \left[ \sum_{k=0}^{\infty} (\rho^2 + \lambda_k^2)^{s/2} (\cos(s\theta_k) + i \sin(s\theta_k)) \right] |\widehat{u_{T,k}}(\rho)|^2 d\rho,
\end{aligned}$$

where  $\theta_k = \tan^{-1}(\frac{\rho}{\lambda_k})$  for  $k \in \mathbb{N}$ . Note that the sine function is odd, and  $|\widehat{u_{T,k}}(\rho)|^2 = |\widehat{u_{T,k}}(-\rho)|^2$  for any  $\rho \in \mathbb{R}$ , then there holds

$$\int_{\mathbb{R}} \sum_{k=0}^{\infty} (\rho^2 + \lambda_k^2)^{s/2} \sin(s\theta_k) |\widehat{u_{T,k}}(\rho)|^2 d\rho = 0.$$

Meanwhile, since  $(\phi_k)_{k \geq 0}$  is an orthonormal basis in  $L^2(M)$ , we can write  $u_T(x, t) = \sum_{k=0}^{\infty} u_{T,k}(t) \phi_k(x)$  such that  $\|u_T(\cdot, t)\|_{L^2(M)}^2 = \sum_{k=0}^{\infty} |u_{T,k}(t)|^2$ , for  $t \in \mathbb{R}$ , where we used  $\|\phi_k\|_{L^2(M)} = 1$ , for all  $k \geq 0$ . Inserting the above identities into (3.2), we can obtain

$$\begin{aligned}
(3.3) \quad |B_g(u_T, u_T)| &= \int_{\mathbb{R}} \sum_{k=0}^{\infty} (\rho^2 + \lambda_k^2)^{s/2} \cos(s\theta_k) |\widehat{u_{T,k}}(\rho)|^2 d\rho \\
&\geq \int_{\mathbb{R}} \sum_{k=0}^{\infty} (\rho^2 + \lambda_k^2)^{s/2} \cos(s\pi/2) |\widehat{u_{T,k}}(\rho)|^2 d\rho \\
&\geq c_s \|u_T\|_{\mathbf{H}^s(M \times \mathbb{R})}^2,
\end{aligned}$$

for some constant  $c_s > 0$  depending only on  $s \in (0, 1)$ .

Therefore, for any  $f \in C_c^\infty(M \times (-T, T))$ , by using (3.1) and (3.3), the Lax-Milgram theorem yields that

$$B_g(u_T, v) = (f, v)_{L^2(M \times \mathbb{R})},$$

admits a unique solution  $u_T \in \mathbf{H}^s(M \times \mathbb{R})$  to the initial value problem (1.1) with  $u_T = 0$  in  $M \times \{t \leq -T\}$ . This proves the assertion.  $\square$

**Remark 3.2.** *By the above well-posedness, the local source-to-solution map (1.3) is well-defined. Moreover, since the solution  $u = 0$  in  $\{t \leq -T\}$  and  $u_T = u$  in  $M \times [-T, T]$ , then it is possible to extend the measurement in  $\mathcal{O} \times \mathbb{R}$ .*

#### 4. THE CALDERÓN PROBLEM

**4.1. Auxiliary lemmas.** It is different to the nonlocal elliptic case as in [FGKU24], the function  $e^{-\tau \mathcal{H}} u$  is not a solution to any (local) "parabolic equations", but we can still find a useful equation for the function  $e^{-\tau \mathcal{H}} u$ . The next lemma is similar to [LLU22, Lemma 3.1] in the Euclidean case.

**Lemma 4.1.** *Let  $u \in \mathbf{H}^s(M \times \mathbb{R})$ , and consider the function*

$$\tilde{u}(x, t, \tau) := e^{-\tau \mathcal{H}} u(x, t), \text{ in } M \times \mathbb{R} \times (0, \infty).$$

*Then  $\tilde{u}$  is a solution to*

$$(4.1) \quad \begin{cases} (\partial_t + \partial_\tau - \Delta_g) \tilde{u} = 0 & \text{in } M \times \mathbb{R} \times (0, \infty), \\ \tilde{u}(x, t, 0) = u(x, t) & \text{in } M \times \mathbb{R}. \end{cases}$$

*Moreover,  $\tilde{u}(x, -T, \tau) = 0$  for any  $x \in M$  and  $\tau \in (0, \infty)$ .*



*Proof.* By using the representation formula (2.7) of  $\tilde{u}(x, t, \tau)$ , we have

$$\tilde{u}(x, t, \tau) = \int_M e^{-\tau\mathcal{L}}(x, z)u(z, t - \tau) dz,$$

so that

$$\begin{aligned} (\partial_\tau - \Delta_g)\tilde{u} &= \int_M \underbrace{[(\partial_\tau - \Delta_g)e^{-\tau\mathcal{L}}(x, z)]}_{=0 \text{ since } e^{-\tau\mathcal{L}}(x, z) \text{ is the heat kernel}} u(t - \tau, z) dz \\ (4.2) \quad &+ \int_M e^{-\tau\mathcal{L}}(x, z)\partial_\tau(u(z, t - \tau)) dz \\ &= \int_M e^{-\tau\mathcal{L}}(x, z)\partial_\tau(u(t - \tau, z)) dz, \end{aligned}$$

for  $(x, t, \tau) \in M \times \mathbb{R} \times (0, \infty)$ , where we used that  $e^{-\tau\mathcal{L}}(x, z)$  is the heat kernel of  $\partial_\tau - \Delta_g$  on  $M$ . Interchanging  $\partial_\tau$  and  $\partial_t$  in the right hand side of (4.2), one has

$$(\partial_\tau - \Delta_g)\tilde{u} = -\partial_t \left( \int_M e^{-\tau\mathcal{L}}(x, z)u(z, t - \tau) dz \right) = -\partial_t \tilde{u} \text{ in } M \times \mathbb{R} \times (0, \infty),$$

which shows the first equation of (4.1) holds. Moreover, it is easy to check that

$$\tilde{u}(x, t, 0) = \lim_{\tau \rightarrow 0} \int_M e^{-\tau\mathcal{L}}(x, z)u(z, t - \tau) dz = u(x, t), \text{ for } (x, t) \in M \times \mathbb{R}.$$

Now, given any  $\tau > 0$ , we can directly see that

$$\tilde{u}(x, -T, \tau) = \int_M e^{-\tau\mathcal{L}}(x, z) \underbrace{u_T(z, -T - \tau)}_{=0 \text{ for } \tau > 0} dz = 0, \text{ for } (x, \tau) \in M \times (0, \infty).$$

This concludes the proof.  $\square$

Let us reformulate the initial value problem (1.1) into a source problem, which helps us eliminate the initial condition in (1.1). As we discussed in Section 3, the solution  $u$  of (1.1) in  $\Omega_T$  is the same as the solution  $u_T(x, t) = \chi_{(-T, T)}(t)u(x, t)$ . Moreover, with the representation formula (2.10) and the pass time condition in (1.1) at hand, we can compute that

$$(4.3) \quad \mathcal{H}^s u(x, t)|_{t \leq -T} = \int_0^\infty \int_M \mathcal{K}_s(x, z; \tau) (u(z, t - \tau) - u(x, t)) dz d\tau \Big|_{\{t \leq -T\}} = 0 \text{ in } M,$$

since  $u(x, t) = u(z, t - \tau) = 0$  for  $t \leq -T$  and for all  $\tau \in (0, \infty)$ . This also shows that the condition (1.2) must be satisfied. In Lemma 2.4, we have explained that the future data of  $u|_{\{t \geq T\}}$  will not affect the solution  $u$  of (1.1) in the spacetime domain  $M \times (-T, T)$ . Hence, there would be infinitely many functions defined globally on  $M \times \mathbb{R}$  solving the equation (1.1) by varying the function  $u|_{\{t \geq T\}}$ . However, combining with (4.3), let us adopt the notion for the "uniqueness" that (1.1) admits a unique solution  $u_T \in \mathbf{H}_{M \times [-T, T]}^s$ . In other words, the equation (1.1) can be rewritten in terms of

$$(4.4) \quad \mathcal{H}^s u_T = f \text{ in } M \times (-T, T),$$

for any  $f \in C_c^\infty(\mathcal{O} \times (-T, T))$ .

By using the previous discussions, we can write the solution  $u_T$  in (4.4) of the form

$$u_T = \mathcal{H}^{-s} f \text{ in } M \times (-T, T),$$

and we aim to write down a more explicit expression formula of  $\mathcal{H}^{-s}$ .

**Lemma 4.2.** *Given  $0 < s < 1$ , let  $f = f(x, t) \in C_c^\infty(M \times (-T, T))$ , then the solution  $u_T$  of (4.4) is given by*

$$(4.5) \quad u_T = \frac{1}{\Gamma(s)} \int_0^\infty e^{-\tau \mathcal{H}} f \frac{d\tau}{\tau^{1-s}} = \int_0^\infty \int_M \mathcal{K}_{-s}(x, z; \tau) f(z, t - \tau) dz d\tau,$$

where  $\mathcal{K}_{-s}(x, z, \tau)$  is the kernel function defined by (2.11) whenever the exponent  $s$  is replaced by  $-s$ .

*Proof.* The proof can be seen by using the formula of the Gamma function that

$$(i\rho + \lambda_k)^{-s} = \frac{1}{\Gamma(s)} \int_0^\infty e^{-\tau(i\rho + \lambda_k)} \frac{d\tau}{\tau^{1-s}}.$$

The derivation in the first identity of (4.5) is similar to the proof of Lemma 2.1. For the second equality in (4.5), one simply uses the heat kernel  $e^{-\tau \mathcal{L}}(x, z)$  combining with the factor  $\frac{1}{\Gamma(s)} \frac{1}{\tau^{1-s}}$ . This proves the assertion.  $\square$

**Remark 4.3.** *The first identity of (4.5) will play an essential role in the proof of Theorem 1.1.*

**4.2. Proof of Theorem 1.1.** Now, we can prove Theorem 1.1.

*Proof of Theorem 1.1.* Let  $\omega_1, \omega_2 \subset \mathcal{O}$  be nonempty open sets such that  $\overline{\omega_1} \cap \overline{\omega_2} = \emptyset$ . Given an arbitrary  $f \in C_c^\infty(\omega_1 \times (-T, T))$ , the condition (1.6) implies that

$$(\partial_t - \Delta_{g_1})^m f = (\partial_t - \Delta_{g_2})^m f = (\partial_t - \Delta_g)^m f \text{ on } \omega_1 \times (-T, T),$$

and  $\Delta_g^m f \in C_c^\infty(\omega_1 \times (-T, T))$ , for any  $m \in \mathbb{N}$ . Meanwhile, using (1.7), there holds that

$$(4.6) \quad \mathcal{H}_1^{-s} ((\partial_t - \Delta_g)^m f)|_{\mathcal{O} \times (-T, T)} = \mathcal{H}_2^{-s} ((\partial_t - \Delta_g)^m f)|_{\mathcal{O} \times (-T, T)},$$

for any  $m \in \mathbb{N}$ , where  $\mathcal{H}_j := \partial_t - \Delta_{g_j}$ , for  $j = 1, 2$ . Note that

$$w_j(x, t) := \chi_{[-T, T]}(t) \mathcal{H}_j^{-s} ((\partial_t - \Delta_{g_j})^m f)(x, t)$$

is the solution to

$$\mathcal{H}_j^s w_j = (\partial_t - \Delta_g)^m f \in C_c^\infty(\omega_1 \times (-T, T)) \quad \text{in } M \times \mathbb{R},$$

for  $j = 1, 2$  and for any  $m \in \mathbb{N}$ . Then the identity (4.6) is equivalent to

$$\mathcal{H}_1^{-s} ((\partial_t - \Delta_g)^m f)|_{\mathcal{O} \times \mathbb{R}} = \mathcal{H}_2^{-s} ((\partial_t - \Delta_g)^m f)|_{\mathcal{O} \times \mathbb{R}},$$

where we use the same technique by multiplying the time cutoff function  $\chi_{[-T, T]}(t)$ , which was shown in Section 3. This makes that the solution  $w_j$  are supported in  $M \times [-T, T]$ , which are zero in the sets  $M \times (\{t \leq -T\} \cup \{t \geq T\})$ , for  $j = 1, 2$ . Let us emphasize again that the future information of these solutions in the set  $M \times \{t \geq T\}$  will not affect the solution in the set  $M \times (-T, T)$ , and hence we simply take the solution to be zero in the future time domain.

By using Lemma 4.2, the above identity yields that

$$(4.7) \quad \int_0^\infty (e^{-\tau \mathcal{H}_1} - e^{-\tau \mathcal{H}_2}) ((\partial_t - \Delta_g)^m f) \frac{d\tau}{\tau^{1-s}} = 0,$$

for  $(x, t) \in \mathcal{O} \times \mathbb{R}$  and  $m \in \mathbb{N}$ . Furthermore, for  $f \in C_c^\infty(\omega_1 \times (-T, T))$  and  $\frac{n+1}{2} < \ell \in \mathbb{N}$ , the Sobolev embedding and (2.8) imply

$$(4.8) \quad \begin{aligned} & \|e^{-\tau \mathcal{H}_j} (\partial_t - \Delta_g)^m f\|_{L^\infty(M_j \times \mathbb{R})} \\ & \leq C \|e^{-\tau \mathcal{H}_j} (\partial_t - \Delta_g)^m f\|_{H^\ell(M_j \times \mathbb{R})} \\ & \leq C \|e^{-\tau \mathcal{H}_j} (1 - \partial_t^2 - \Delta_{g_j})^\ell (\partial_t - \Delta_g)^m f\|_{L^2(M_j \times \mathbb{R})} \\ & \leq C e^{-c_0 \tau} \| (1 - \partial_t^2 - \Delta_{g_j})^\ell (\partial_t - \Delta_g)^m f \|_{L^2(M_j \times \mathbb{R})} \end{aligned}$$

which ensures that the the integral in (4.7) convergence uniformly for  $(x, t) \in \mathcal{O} \times \mathbb{R}$ , for any  $m \in \mathbb{N}$ . Notice that in the right-hand side of (4.8), it is a local computation, for any smooth function  $f \in C_c^\infty(\omega_1 \times (-T, T))$ .

Similar to [LLU22], using the commutative property  $e^{-\tau \mathcal{H}_j} \mathcal{H}_j^m = \mathcal{H}_j^m e^{-\tau \mathcal{H}_j}$ , for all  $\tau \geq 0$  on  $\text{Dom}(\mathcal{H}_j^m)$ , we can obtain

$$(4.9) \quad e^{-\tau \mathcal{H}_j} ((\partial_t - \Delta_g)^m f) = (\partial_t - \Delta_g)^m (e^{-\tau \mathcal{H}_j} f) = \underbrace{(-\partial_\tau)^m (e^{-\tau \mathcal{H}_j} f)}_{\text{By (4.1)}},$$

for any  $m \in \mathbb{N}$ , where  $e^{-\tau \mathcal{H}_j} f$  solves (4.1) as  $g = g_j$ , for  $j = 1, 2$ . Inserting (4.9) into (4.7), there holds that

$$(4.10) \quad \int_0^\infty \partial_\tau^m ((e^{-\tau \mathcal{H}_1} - e^{-\tau \mathcal{H}_2}) f) (x, t) \frac{d\tau}{\tau^{1-s}} = 0,$$

for  $(x, t) \in \mathcal{O} \times \mathbb{R}$  and  $m \in \mathbb{N}$ . Meanwhile, as  $(x, t) \in \omega_2 \times (-T, T)$ , we can integrate by parts (4.10)  $m$ -times. Note that there are no end-point contributions while doing this integration by parts. More concretely, as showed in [FGKU24], using the exponential decay (4.8) as  $\tau \rightarrow \infty$  and the Gaussian upper bound (2.5) for the heat kernel  $e^{-\tau \mathcal{L}}(x, z)$  as  $\tau \rightarrow 0$ , one can guarantee that

$$(4.11) \quad \begin{aligned} & \lim_{\tau \rightarrow \infty} \left| \partial_\tau^\ell ((e^{-\tau \mathcal{H}_1} - e^{-\tau \mathcal{H}_2}) f) (x, t) \frac{1}{\tau^{k-s}} \right| \\ &= \lim_{\tau \rightarrow 0} \left| \partial_\tau^\ell ((e^{-\tau \mathcal{H}_1} - e^{-\tau \mathcal{H}_2}) f) (x, t) \frac{1}{\tau^{k-s}} \right| = 0, \end{aligned}$$

where we use

$$(4.12) \quad \left| \partial_\tau^\ell ((e^{-\tau \mathcal{H}_1} - e^{-\tau \mathcal{H}_2}) f) (x, t) \right| \leq C e^{-\frac{c}{\tau}}$$

for any  $k, \ell \in \mathbb{N}$ , for any  $\tau \in (0, 1)$  and for some constant  $c > 0$ .

With (4.11) at hand, integrating (4.10)  $m$ -times, there holds that

$$\int_0^\infty ((e^{-\tau \mathcal{H}_1} - e^{-\tau \mathcal{H}_2}) f) (x, t) \frac{d\tau}{\tau^{m+1-s}} = 0, \text{ for all } m \in \mathbb{N},$$

which is equivalent to

$$(4.13) \quad \int_0^\infty ((e^{-\tau \mathcal{H}_1} - e^{-\tau \mathcal{H}_2}) f) (x, t) \frac{d\tau}{\tau^{m+2-s}} = 0, \text{ for all } m \in \mathbb{N} \cup \{0\}.$$

By using the change of variable that  $\eta = \frac{1}{\tau}$ , the identity (4.13) yields that

$$(4.14) \quad \int_0^\infty \varphi(\eta) \eta^m d\eta = 0,$$

for all  $m \in \mathbb{N}$ , where

$$\varphi(\eta) := \frac{((e^{-\frac{1}{\eta} \mathcal{H}_1} - e^{-\frac{1}{\eta} \mathcal{H}_2}) f) (x, t)}{\eta^s}, \text{ for } (x, t) \in \omega_2 \times \mathbb{R}.$$

By using (2.8) and (4.12), one can obtain that

$$|\varphi(\eta)| \leq C \frac{e^{-c\eta}}{\eta^s},$$

for some constants  $c, C > 0$ . Therefore, the Fourier transform of  $\chi_{[0, \infty)}(\eta) \varphi(\eta)$  is

$$\mathcal{F}(\chi_{[0, \infty)}(\eta) \varphi(\eta))(\xi) = \frac{1}{\sqrt{2\pi}} \int_0^\infty \varphi(\eta) e^{-i\eta \xi} d\eta$$

is holomorphic for  $\text{Im}\xi < c$ . Via (4.14), it is known that  $\mathcal{F}(\chi_{[0,\infty)}\varphi)(\xi)$  equals to zero at  $\xi = 0$ , which implies that  $\varphi(\eta) \equiv 0$  for  $\eta > 0$ . This implies that

$$(4.15) \quad ((e^{-\tau\mathcal{H}_1} - e^{-\tau\mathcal{H}_2})f)(x, t) = 0 \text{ for } (x, t) \in \omega_2 \times \mathbb{R} \text{ and } \tau > 0.$$

The next task is to transfer the above identity into another useful identity (1.8) as we mentioned.

Thanks to Lemma 4.1, we know that  $e^{-\tau\mathcal{H}_j}f$  solves the equation

$$(\partial_\tau - \Delta_{g_j})(e^{-\tau\mathcal{H}_j}f) = -\partial_t(e^{-\tau\mathcal{H}_j}f), \text{ for } (x, t, \tau) \in M \times \mathbb{R} \times (0, \infty),$$

for  $j = 1, 2$ . Let us integrate the above equation with respect to  $t \in \mathbb{R}$ , so that

$$(4.16) \quad \begin{aligned} (\partial_\tau - \Delta_{g_j})F_j &= -\int_{-\infty}^{\infty} \partial_t(e^{-\tau\mathcal{H}_j}f) dt \\ &= \lim_{t \rightarrow -\infty} \left( \int_M e^{-\tau\mathcal{L}_j}(x, z)f(z, t - \tau) dz \right) \\ &\quad - \lim_{t \rightarrow \infty} \left( \int_M e^{-\tau\mathcal{L}_j}(x, z)f(z, t - \tau) dz \right) \\ &= 0, \end{aligned}$$

for all  $x \in M$ ,  $\tau > 0$  and  $f \in C_c^\infty(\omega_1 \times (-T, T))$ , where  $\mathcal{L}_j := -\Delta_{g_j}$ , and  $F_j = \int_{-\infty}^{\infty} (e^{-\tau\mathcal{H}_j}f) dt$ , for  $j = 1, 2$ .

In addition, observe the function  $F_j(x, \tau)$  more carefully, one can rewrite  $F_j$  in terms of

$$(4.17) \quad \begin{aligned} F_j(x, \tau) &= \int_{-\infty}^{\infty} (e^{-\tau\mathcal{H}_j}f)(x, t) dt \\ &= \int_{-\infty}^{\infty} \int_M e^{-\tau\mathcal{L}_j}(x, z)f(z, t - \tau) dz dt \\ &= \underbrace{\int_M e^{-\tau\mathcal{L}_j}(x, z) \left( \int_{-\infty}^{\infty} f(z, t - \tau) dt \right) dz}_{\text{By the Fubini (Tonelli) theorem}} \\ &= (e^{-\tau\mathcal{L}_j}\mathbf{f})(x), \end{aligned}$$

for  $j = 1, 2$ , which are the well-defined, since the right-hand side in (4.17) is the heat kernel representation formula as  $\mathbf{f} \in C_c^\infty(\omega_1)$ . Here  $e^{-\tau\mathcal{L}_j}(x, z) \geq 0$  is the heat kernel of  $\partial_\tau + \mathcal{L}_j$  for  $j = 1, 2$ , and

$$\mathbf{f}(z) := \int_{-\infty}^{\infty} f(z, t - \tau) dt \in C_c^\infty(\omega_1),$$

since  $f \in C_c^\infty(\omega_1 \times (-T, T))$ . Due to arbitrariness of  $f(z, t - \tau) \in C_c^\infty(\omega_1 \times (-T, T))$ , the function  $\mathbf{f}(z) \in C_c^\infty(\omega_1)$  can be also arbitrary<sup>1</sup>.

Now, on the one hand, integrate (4.15) with respect to  $t \in (-\infty, \infty)$ , plugging (4.17) into (4.15) (after  $t$ -integration), then there holds that

$$((e^{-\tau\mathcal{L}_1} - e^{-\tau\mathcal{L}_2})\mathbf{f})(x) = 0 \text{ for } x \in \omega_2 \text{ and } \tau > 0.$$

On the other hand, without loss of generality, we may assume that  $\mathcal{O}$  is connected, which is contained in a single coordinate patch for both manifolds  $M_1$  and  $M_2$ . Note that the function  $e^{-\tau\mathcal{L}_j}\mathbf{f}$  solves the heat equation (4.16), for  $j = 1, 2$ , which implies that  $(e^{-\tau\mathcal{L}_1} - e^{-\tau\mathcal{L}_2})\mathbf{f}$  solves the heat equation

$$(\partial_\tau - \Delta_g)((e^{-\tau\mathcal{L}_1} - e^{-\tau\mathcal{L}_2})\mathbf{f}) = 0 \text{ for } (x, \tau) \in \mathcal{O} \times (0, \infty).$$

<sup>1</sup>This can be easily seen by the following simple observations: Given any  $\mathbf{f} \in C_c^\infty(\omega_1)$ , choosing  $f(z, t - \tau) := \mathbf{f}(z)h(t - \tau)$ , where  $h(\cdot) \in C_c^\infty(-T, T)$  such that  $\int_{-\infty}^{\infty} h(t) dt = \int_{-\infty}^{\infty} h(t - \tau) dt = 1$ , for all  $\tau > 0$ .

Then the unique continuation property of the heat equation (cf. [Lin90]) implies

$$\left( (e^{-\tau\mathcal{L}_1} - e^{-\tau\mathcal{L}_2}) \mathbf{f} \right) = 0 \text{ for } (x, \tau) \in \mathcal{O} \times (0, \infty).$$

Now, as  $f(x, t) \in C_c^\infty(\omega_1 \times (-T, T))$  and  $\omega_1 \Subset \mathcal{O}$  arbitrary, so  $\mathbf{f}(x) \in C_c^\infty(\omega_1)$  and  $\omega_1$  can be also arbitrary.

Therefore, one can obtain

$$e^{-\tau\mathcal{L}_1} \mathbf{f}|_{\mathcal{O}} = e^{-\tau\mathcal{L}_2} \mathbf{f}|_{\mathcal{O}}, \text{ for } \tau > 0.$$

Due to the arbitrariness of  $\mathbf{f} \in C_c^\infty(\mathcal{O})$ , the above identity implies

$$e^{-\tau\mathcal{L}_1}(x, z) = e^{-\tau\mathcal{L}_2}(x, z), \text{ for } x, z \in \mathcal{O} \text{ and } \tau > 0,$$

i.e., (1.8) holds. Finally, applying [FGKU24, Theorem 1.5], one can find a diffeomorphism  $\Phi : M_1 \rightarrow M_2$  such that  $\Phi^*g_2 = g_1$  on  $M_1$  with  $\Phi = \text{Id}$  on  $\mathcal{O}$ . This proves the assertion.  $\square$

**Acknowledgments.** The author is partially supported by the National Science and Technology Council (NSTC) Taiwan, under the projects 113-2628-M-A49-003 & 113-2115-M-A49-017-MY3. The author is also a Humboldt research fellow (for experienced researchers) in Germany.

#### REFERENCES

- [BDLCS21] Animesh Biswas, Marta De León-Contreras, and Pablo Raúl Stinga. Harnack inequalities and Hölder estimates for master equations. *SIAM Journal on Mathematical Analysis*, 53(2):2319–2348, 2021.
- [BS24] Agnid Banerjee and Soumen Senapati. The Calderón problem for space-time fractional parabolic operators with variable coefficients. *SIAM J. Math. Anal.*, 56(4):4759–4810, 2024.
- [Cal06] Alberto P. Calderón. On an inverse boundary value problem. *Comput. Appl. Math.*, 25(2-3):133–138, 2006.
- [CGRU23] Giovanni Covi, Tuhin Ghosh, Angkana Rüland, and Gunther Uhlmann. A reduction of the fractional Calderón problem to the local Calderón problem by means of the Caffarelli-Silvestre extension. *arXiv preprint arXiv:2305.04227*, 2023.
- [CLL19] Xinlin Cao, Yi-Hsuan Lin, and Hongyu Liu. Simultaneously recovering potentials and embedded obstacles for anisotropic fractional Schrödinger operators. *Inverse Probl. Imaging*, 13(1):197–210, 2019.
- [CLR20] Mihajlo Cekic, Yi-Hsuan Lin, and Angkana Rüland. The Calderón problem for the fractional Schrödinger equation with drift. *Cal. Var. Partial Differential Equations*, 59(91), 2020.
- [CMR21] Giovanni Covi, Keijo Mönkkönen, and Jesse Railo. Unique continuation property and Poincaré inequality for higher order fractional Laplacians with applications in inverse problems. *Inverse Probl. Imaging*, 15(4):641–681, 2021.
- [CMRU22] Giovanni Covi, Keijo Mönkkönen, Jesse Railo, and Gunther Uhlmann. The higher order fractional Calderón problem for linear local operators: Uniqueness. *Adv. Math.*, 399:Paper No. 108246, 2022.
- [Fei24] Ali Feizmohammadi. Fractional Calderón problem on a closed Riemannian manifold. *Trans. Amer. Math. Soc.*, 377(4):2991–3013, 2024.
- [FGKU24] Ali Feizmohammadi, Tuhin Ghosh, Katya Krupchyk, and Gunther Uhlmann. Fractional anisotropic Calderón problem on closed Riemannian manifolds. *J. Diff. Geom.*, to appear, 2024.
- [FKU24] Ali Feizmohammadi, Katya Krupchyk, and Gunther Uhlmann. Calderón problem for fractional Schrödinger operators on closed Riemannian manifolds. *arXiv preprint arXiv:2407.16866*, 2024.
- [FL24] Ali Feizmohammadi and Yi-Hsuan Lin. Entanglement principle for the fractional Laplacian with applications to inverse problems. *arXiv preprint arXiv:2412.13118*, 2024.
- [GLX17] Tuhin Ghosh, Yi-Hsuan Lin, and Jingni Xiao. The Calderón problem for variable coefficients nonlocal elliptic operators. *Comm. Partial Differential Equations*, 42(12):1923–1961, 2017.

- [GRSU20] Tuhin Ghosh, Angkana Rüland, Mikko Salo, and Gunther Uhlmann. Uniqueness and reconstruction for the fractional Calderón problem with a single measurement. *J. Funct. Anal.*, 279(1):108505, 42, 2020.
- [GSU20] Tuhin Ghosh, Mikko Salo, and Gunther Uhlmann. The Calderón problem for the fractional Schrödinger equation. *Anal. PDE*, 13(2):455–475, 2020.
- [GU21] Tuhin Ghosh and Gunther Uhlmann. The Calderón problem for nonlocal operators. *arXiv:2110.09265*, 2021.
- [HL19] Bastian Harrach and Yi-Hsuan Lin. Monotonicity-based inversion of the fractional Schrödinger equation I. Positive potentials. *SIAM J. Math. Anal.*, 51(4):3092–3111, 2019.
- [HL20] Bastian Harrach and Yi-Hsuan Lin. Monotonicity-based inversion of the fractional Schrödinger equation II. General potentials and stability. *SIAM J. Math. Anal.*, 52(1):402–436, 2020.
- [KLW22] Pu-Zhao Kow, Yi-Hsuan Lin, and Jenn-Nan Wang. The Calderón problem for the fractional wave equation: uniqueness and optimal stability. *SIAM J. Math. Anal.*, 54(3):3379–3419, 2022.
- [KLZ24] Manas Kar, Yi-Hsuan Lin, and Philipp Zimmermann. Determining coefficients for a fractional  $p$ -Laplace equation from exterior measurements. *J. Differential Equations*, 406:338–365, 2024.
- [KRZ23] Manas Kar, Jesse Railo, and Philipp Zimmermann. The fractional  $p$ -biharmonic systems: optimal Poincaré constants, unique continuation and inverse problems. *Calc. Var. Partial Differential Equations*, 62(4):Paper No. 130, 36, 2023.
- [Lin90] Fang-Hua Lin. A uniqueness theorem for parabolic equations. *Comm. Pure Appl. Math.*, 43(1):127–136, 1990.
- [Lin22] Yi-Hsuan Lin. Monotonicity-based inversion of fractional semilinear elliptic equations with power type nonlinearities. *Calc. Var. Partial Differential Equations*, 61(5):Paper No. 188, 30, 2022.
- [Lin23] Yi-Hsuan Lin. Determining both leading coefficient and source in a nonlocal elliptic equation. *arXiv preprint arXiv:2312.15607*, 2023.
- [LL22] Ru-Yu Lai and Yi-Hsuan Lin. Inverse problems for fractional semilinear elliptic equations. *Nonlinear Anal.*, 216:Paper No. 112699, 21, 2022.
- [LL23] Yi-Hsuan Lin and Hongyu Liu. Inverse problems for fractional equations with a minimal number of measurements. *Communications and Computational Analysis*, 1:72–93, 2023.
- [LLR20] Ru-Yu Lai, Yi-Hsuan Lin, and Angkana Rüland. The Calderón problem for a space-time fractional parabolic equation. *SIAM J. Math. Anal.*, 52(3):2655–2688, 2020.
- [LLS20] Matti Lassas, Tony Liimatainen, and Mikko Salo. The Poisson embedding approach to the Calderón problem. *Math. Ann.*, 377(1-2):19–67, 2020.
- [LLU22] Ching-Lung Lin, Yi-Hsuan Lin, and Gunther Uhlmann. The Calderón problem for nonlocal parabolic operators. *arXiv preprint arXiv:2209.11157*, 2022.
- [LLU23] Ching-Lung Lin, Yi-Hsuan Lin, and Gunther Uhlmann. The Calderón problem for nonlocal parabolic operators: A new reduction from the nonlocal to the local. *arXiv preprint arXiv:2308.09654*, 2023.
- [LNZ24] Yi-Hsuan Lin, Gen Nakamura, and Philipp Zimmermann. The Calderón problem for the Schrödinger equation in transversally anisotropic geometries with partial data. *arXiv preprint arXiv:2408.08298*, 2024.
- [LTZ24a] Yi-Hsuan Lin, Teemu Tyni, and Philipp Zimmermann. Optimal Runge approximation for nonlocal wave equations and unique determination of polyhomogeneous nonlinearities. *arXiv preprint arXiv:2408.13869*, 2024.
- [LTZ24b] Yi-Hsuan Lin, Teemu Tyni, and Philipp Zimmermann. Well-posedness and inverse problems for semilinear nonlocal wave equations. *Nonlinear Anal.*, 247:Paper No. 113601, 19, 2024.
- [LU89] John M. Lee and Gunther Uhlmann. Determining anisotropic real-analytic conductivities by boundary measurements. *Comm. Pure Appl. Math.*, 42(8):1097–1112, 1989.
- [LU01] Matti Lassas and Gunther Uhlmann. On determining a Riemannian manifold from the Dirichlet-to-Neumann map. *Ann. Sci. École Norm. Sup. (4)*, 34(5):771–787, 2001.
- [LZ23] Yi-Hsuan Lin and Philipp Zimmermann. Unique determination of coefficients and kernel in nonlocal porous medium equations with absorption term. *arXiv preprint arXiv:2305.16282*, 2023.
- [LZ24] Yi-Hsuan Lin and Philipp Zimmermann. Approximation and uniqueness results for the nonlocal diffuse optical tomography problem. *arXiv preprint arXiv:2406.06226*, 2024.

- [RS18] Angkana Rüland and Mikko Salo. Exponential instability in the fractional Calderón problem. *Inverse Problems*, 34(4):045003, 21, 2018.
- [RS20] Angkana Rüland and Mikko Salo. The fractional Calderón problem: low regularity and stability. *Nonlinear Anal.*, 193:111529, 56, 2020.
- [Rül23] Angkana Rüland. Revisiting the anisotropic fractional Calderón problem using the Caffarelli-Silvestre extension. *arXiv preprint arXiv:2309.00858*, 2023.
- [Sam02] Stefan G. Samko. *Hypersingular integrals and their applications*, volume 5 of *Analytical Methods and Special Functions*. Taylor & Francis Group, London, 2002.
- [SY94] Richard M Schoen and Shing-Tung Yau. *Lectures on differential geometry*, volume I of *Conference Proceedings and Lecture Notes in Geometry and Topology*. International Press, Cambridge, MA, 1994. Lecture notes prepared by Wei Yue Ding, Kung Ching Chang [Gong Qing Zhang], Jia Qing Zhong and Yi Chao Xu, Translated from the Chinese by Ding and S. Y. Cheng, With a preface translated from the Chinese by Kaising Tso.

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL YANG MING CHIAO TUNG UNIVERSITY, HSINCHU, TAIWAN & FAKULTÄT FÜR MATHEMATIK, UNIVERSITY OF DUISBURG-ESSEN, ESSEN, GERMANY

*Email address:* [yihuanlin3@gmail.com](mailto:yihuanlin3@gmail.com)